National Library of Canada
Bibliothèque nationale du Canada

Canadian Theses Service
Service des thèses canadiennes

Ottawa, Canada
K1A 0N4

NOTICE

The quality of this microform is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Reproduction in full or in part of this microform is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30, and subsequent amendments.

AVIS

La qualité de cette microforme dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

La reproduction, même partielle, de cette microforme est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30, et ses amendements subséquents.

NL-339 (r. 88/04) c

Canada

UNIVERSITY OF ALBERTA

CALCIUM SUPPLEMENTATION IN PREMATURE INFANTS FED BREAST MILK: The Effect of Timing of Calcium Supplementation on Calcium, Phosphorus, Magnesium and Fat Balance and on the Absorption Coefficients of Individual Fatty Acids

BY

Yiping Wang

©

A thesis submitted to the Faculty of Graduate Studies and Research in partial fulfillment of the requirements for the degree of Master of Science.

Department of Foods and Nutrition

Edmonton, Alberta

Spring 1992

National Library of Canada
Bibliothèque nationale du Canada

Canadian Theses Service
Service des thèses canadiennes

Ottawa, Canada
K1A 0N4

The author has granted an irrevocable non-exclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of his/her thesis by any means and in any form or format, making this thesis available to interested persons.

L'auteur a accordé une licence irrévocable et non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de sa thèse de quelque manière et sous quelque forme que ce soit pour mettre des exemplaires de cette thèse à la disposition des personnes intéressées.

The author retains ownership of the copyright in his/her thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without his/her permission.

L'auteur conserve la propriété du droit d'auteur qui protège sa thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

ISBN 0-315-73180-X

Canada

UNIVERSITY OF ALBERTA

RELEASE FORM

NAME OF AUTHOR: Yiping Wang

TITLE OF THESIS: CALCIUM SUPPLEMENTATION IN PREMATURE INFANTS FED BREAST MILK: The Effect of Timing of Calcium Supplementation on Calcium, Phosphorus, Magnesium and Fat Balance and on the Absorption Coefficients of Individual Fatty Acids

DEGREE: Master of Science

YEAR THIS DEGREE GRANTED: 1992

Permission is hereby granted to the University of Alberta Library to reproduce single copies of this thesis and to lend or sell such copies for private, scholarly or scientific research purposes only.

The author reserves all other publication and other rights in association with the copyright in the thesis, and except as hereinbefore provided neither the thesis nor any substantial portion thereof may be printed or otherwise reproduced in any material form whatever without the author's prior written permission.



11419 80 Av.

Edmonton, Alberta

Canada, T6G 0R6

Feb. 25, 1992

UNIVERSITY OF ALBERTA

FACULTY OF GRADUATE STUDIES AND RESEARCH

The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research for acceptance, a thesis entitled **CALCIUM SUPPLEMENTATION IN PREMATURE INFANTS FED BREAST MILK: The Effect of Timing of Calcium Supplementation on Calcium, Phosphorus, Magnesium and Fat Balance and on the Absorption Coefficients of Individual Fatty Acids** submitted by **Yiping Wang** in partial fulfillment of the requirements for the degree of **Master of Science.**

Dr. M.T. Clandinin (Supervisor)

Dr. J. Van Aerde

Dr. M.L. Garg

Dr. T. Basu (Committee Chairman)

Feb. 25, 1992

TO MY MOTHER FOR HER LOVE AND SUPPORT

ABSTRACT

1. Calcium and fatty acid absorption in premature infants can be affected by calcium soap formation with fatty acids in the intestinal lumen following oral calcium supplementation. Three seventy-two-hour balance studies were conducted in sixteen pair-matched premature infants with a gestational age of 30.8 ± 1.4 weeks and a birthweight of 1329 ± 168 grams. Calcium, phosphorus, magnesium, fat balance and the absorption coefficients of individual fatty acids were measured in infants given calcium supplementation between feedings and in infants given calcium supplementation with feedings of expressed preterm human milk.
2. Calcium and phosphorus content of preterm human milk (PTM) decreased gradually in the first four weeks of life.
3. Calcium lactate (1.3 mmole/Kg/day) administration between feedings significantly decreased phosphorus absorption and retention, magnesium retention, and increased total magnesium loss compared to giving calcium with feedings. Calcium absorption tended to be impaired in infants receiving calcium between feedings, although this was not statistically significant.
4. Total lipid and fatty acid C14:0, C16:0, C18:0 and C18:1(9) absorption was also decreased by calcium supplementation between feedings when compared to supplementation with feedings. However, the differences were not statistically significant. Infants given calcium between feedings seem to have gained less weight compared to infants receiving calcium with feedings, especially in the second

postnatal week, although it was not statistically significant.

5. A significant correlation between calcium intake and calcium loss in stool was observed. In both feeding regimens, fecal lipid was positively correlated with calcium and phosphorus content in the stool. Calcium and phosphorus excretions positively correlated with magnesium in the stool in infants given calcium between feedings. In infants receiving calcium with feedings, fecal calcium and magnesium positively correlated with phosphorus concentration in the stool.

6. The percentages of neutral lipid, calcium salts, free fatty acids and compound lipid in stool were 54.4%, 2.5%, 21.5%, and 21.7% respectively. Neutral lipid excretion increased and free fatty acids decreased gradually during the first four weeks of life, although they were not statistically significant.

7. It is concluded that intraluminal hydrolysis of milk lipid might be less complete when supplementing calcium between feedings. Timing change of calcium supplementation from with feedings to between feedings may also complicate divalent ion interaction within the intestinal lumen and eventually the loss of calcium, phosphorus, magnesium, total fat and certain long-chain saturated fatty acids, which would in turn affect growth in premature infants.

ACKNOWLEDGEMENTS

Many thanks first to my parents, brothers, sister, and friends for their support and encouragement during my study.

I wish to thank my supervisors, Dr. M.T. Clandinin and Dr. J. Van Aerde for their guidance throughout the program. A special thank you to Dr. M.T. Clandinin for his valuable advice, encouragement and patience. His time spent on the editorial supervision of this thesis is also highly appreciated. My sincere thank you is extended to Dr. J. Van Aerde for his time, supervision, and friendship. His provision of study samples, laboratory space and materials is very much appreciated.

I am also grateful to Mr. G. Chan and Dr. Y.K. Goh for their comprehensive technical supervision.

Finally, the financial support from UNICEF and the Medical Research Council of Canada is gratefully acknowledged.

TABLE OF CONTENTS

Page

Chapter I: LITERATURE REVIEW AND RATIONALE

A: Introduction.....1

B: Literature Review.....2

1. Introduction.....2

2. Preterm Human Milk vs Term Human Milk or Formulas.....2

Nutrient Composition of Preterm Human Milk.....2

The Superior Fat Absorption.....3

The Role of Fatty Acids in Growth and Development.....4

The Shortage of Calcium and Phosphorus of Preterm Human Milk.....6

The Evidence of Calcium and Phosphorus Deficiency in Premature Infants and Its Improvement after Dietary Supplementation.....7

3. Factors Influencing Intestinal Fat Absorption.....8

Intraluminal Factors.....8

Dietary Factors.....11

4. Factors Influencing Intestinal Calcium Absorption.....15

The Ratio of Dietary Calcium and Phosphorus Intake.....15

Vitamin D and Other Regulating Factors.....15

Dietary Fat.....19

Others.....19

5. Introduction to Phosphorus Absorption.....20

Page

6. Introduction to Magnesium Absorption.....22

7. Summary.....24

C: Rationale.....26

1. Hypothesis.....27

2. Objectives.....27

D: References.....28

Chapter II: CALCIUM SUPPLEMENTATION IN PREMATURE INFANTS FED BREAST MILK:

The Effect of Timing of Calcium Supplementation on Calcium, Phosphorus, Magnesium and Fat Balance and on the Absorption Coefficients of Individual Fatty Acids

A: Introduction.....41

B: Study Subjects and Methodology.....42

1. Study Design.....42

Study Subject Description.....42

Feeding Regimen.....42

Calcium Supplementation.....43

2. Collection of Preterm Human Milk.....43

3. Collections of Urine and Stool Samples.....43

4. Anthropometric Measurements 44

5. Calcium, Phosphorus, Magnesium, Total Fat and Fatty Acids Measurements 44

6. Statistics Applied.....45

C: Results.....45

1. Composition of Preterm Human Milk.....45

2. Anthropometric Data.....47

3. Balance Studies.....49

Page

Lipid Balance Study.....49

Calcium Balance Study.....49

Phosphorus Balance Study.....52

Magnesium Balance Study.....54

4. Correlation of Calcium Parameters.....56

5. Correlations of Fecal Lipid, Calcium, Phosphorus and Magnesium 56

6. Absorption Coefficients of Individual Fatty Acids.....56

7. Fecal Lipid Fractions.....65

D: Discussion and Conclusions.....75

1. Chronological Changes in Milk Composition.....75

2. Growth Parameters in Anthropometric Study.....75

3. Effects of Calcium Supplementation on Intestinal Fat and Mineral Absorption.....76

Possible Activation of Lipolytic Enzyme(s).....77

Calcium Soap Formation.....78

Interaction of Divalent Cations.....78

4. Absorption Coefficients of Individual Fatty Acids.....79

5. Fecal Lipid Fractions.....79

E: References 82

Appendices: I. Extraction of Milk Lipid.....84

II. Extraction of Fecal Lipid.....85

List of Tables

Page

II-1. Composition of Expressed Preterm Human Milk Consumed by Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....46

II-2. Anthropometric Data for Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....48

II-3. Lipid Balance in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....50

II-4. Calcium Balance in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....51

II-5. Phosphorus Balance in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....53

II-6. Magnesium Balance in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....55

II-7. Absorption Coefficients for Selected Fatty Acids in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....63

II-8. Quantitative Measurement of Selected Fecal Fatty Acids in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....64

II-9. Percentage of Fecal Lipid Fractions in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....66

Page

II-10. Quantitative Measurement of Fecal Lipid Fractions in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....68

II-11a. Quantitative Measurement of Fecal Fatty Acid C12:0 in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week.....69

II-11b. Quantitative Measurement of Fecal Fatty Acid C14:0 in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week.....70

II-11c. Quantitative Measurement of Fecal Fatty Acid C16:0 in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week.....71

II-11d. Quantitative Measurement of Fecal Fatty Acid C18:0 in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week.....72

II-11e. Quantitative Measurement of Fecal Fatty Acid C18:1(9) in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week.....73

II-11f. Quantitative Measurement of Fecal Fatty Acid C18:2(6) in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week.....74

List of Figures

Page

II-1. Correlation of Ca Intake and Ca Loss in Stool in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....57

II-2. Correlation of Fecal Lipid and Ca Losses in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....58

II-3. Correlation of Fecal Lipid and P Losses in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....59

II-4. Correlation of Fecal P and Mg Losses in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week.....60

II-5. Correlation of Fecal Ca and P Losses in Breast Fed Premature Infants Receiving Ca Lactate With Feedings in the First, Second and Fourth Postnatal Week.....61

II-6. Correlation of Fecal Ca and Mg Losses in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings in the First, Second and Fourth Postnatal Week.....62

Chapter I

LITERATURE REVIEW AND RATIONALE

A: Introduction

Preterm human milk has been highly recommended to premature infants for its suitable macronutrient content and immunological protection. However, calcium and phosphorus inadequacy in their mother's own milk put these infants at high risk of developing various nutritional abnormalities and eventually growth retardation. The practice of calcium supplementation in premature infants fed breast milk has been proven to reduce this risk significantly. As increased calcium intake interferes with the absorption of fat and other divalent cations, it is postulated that the change in timing of calcium supplementation from with feedings to between feedings will reduce these interactions. This chapter will review the content of certain nutrients in preterm human milk as compared to either term human milk or a commercial formula, the evidence of calcium and phosphorus deficiency in the premature infant and the effect of calcium supplementation. Factors affecting intestinal fat and calcium absorption will also be discussed, as well as intestinal phosphorus and magnesium absorption. Finally, the rationale, hypothesis and objectives of this thesis will be presented.

B: Literature Review

1. Introduction

Calcium supplementation has been found to relate to increased fecal lipid loss in numerous studies with orally fed premature infants. Following supplementation of calcium lactate the amount of calcium absorbed is significantly increased, but fecal excretion of fat increases as well, resulting in a significantly lower percentage of ingested fat being absorbed in association with calcium supplementation (1). This complication and the more appropriate calcium supplementation require further study.

2. Preterm Human Milk vs Term Human Milk or Formulas

Nutrient Compositions of Preterm Human Milk

Mother's own breast milk has been increasingly recognized as the optimal food for premature infants as a source of nutrition in the first weeks of life. When premature infants are given a mixture of fresh human milk and an infant formula, fat absorption is greater than for infants receiving the formula alone (2). In addition, comparing to banked mature human milk or infant formula, the protein and amino acid content of preterm human milk offers about 75% more protein relative to energy, enabling growth of the premature infant to approach intrauterine nitrogen retention rates. Thus a greater net weight gain is achieved at least in the first two weeks of life (3,4,5). Furthermore, lactose present in

human milk can be easily digested and absorbed by neonates, and improves calcium absorption as well (6). Therefore, in order to meet the estimated macronutrient requirement for premature infants to reach an intrauterine growth rate, preterm human milk is more appropriate than mature human milk during the first weeks of life.

The Superior Fat Absorption of Preterm Human Milk

The composition of preterm milk enables superior fat absorption. First, preterm milk contains more medium-chain fatty acids and long-chain polyunsaturated fatty acids that have better absorption coefficients than full-term milk (7). Second, a small amount of bile salt-stimulated lipase is present in human milk, which, to a certain extent, may compensate for the shortage of pancreatic lipase and bile salts in the premature infant and help to hydrolyze fat in intestinal lumen (7). Because this enzyme is relatively heat-unstable, premature infants fed pooled pasteurized preterm milk tend to grow more slowly than those fed fresh preterm milk (8). Pancreatic lipase in the intestinal lumen preferably hydrolyzes triglycerides to beta-monoglyceride, while the bile salt-stimulated lipase in human milk does not have this positional specificity. Therefore, in infants fed human milk, beta-monoglyceride produced by pancreatic lipase is well hydrolyzed by this bile salt-stimulated lipase and thus the intestinal hydrolysis of milk lipid is relatively complete (9). Because of superior fat absorption, premature infants fed their mothers' own milk gain more weight when compared to either formula or pooled and heat-treated breast milk (3,10).

The Role of Fatty Acids in Growth and Development

Human milk contains long-chain polyunsaturated fatty acids, whereas most infant formulas lack this essential element (11). Because a significant amount of omega-3 fatty acids, especially docosahexaenoic acid (DHA), accumulates in the adipose tissue during the last trimester of pregnancy mainly through placental transfer, and preferential desaturation of long-chain omega-3 fatty acids in brain tissue occurs only after 30 weeks gestation, provision of premature infants with these fatty acids is critically important both for energy production and for synthesis of structural lipid (12,13,14).

Omega-3 long-chain polyunsaturated fatty acids are especially important in the growth and development of eyes and brain of neonates, as animal studies show that dietary omega-3 fatty acid deficiency during pregnancy and lactation affects normal vision, causes irreversible changes in electroretinograms and impairs learning ability in the offspring (15,16). It is suggested that omega-3 fatty acids in the membranes of photorecepter cells and synapses are important in biogenesis of these membranes during the perinatal period, in normal function of the tissues and in response of the nervous system to ischemia, convulsion and retinal stimulation (17). In rats, omega-3 fatty acid deficiency increases the brain's susceptibility to exogeneous toxins (18). An altered fatty acid profile of cerebroside in phospholipid in developing brain may alter myelination as well (19).

As the proportion of omega-3 fatty acids increases gradually only after 30 week gestation through the maturation of the chain elongation and

desaturation enzymatic system (14), nutrition during the first weeks of life in premature infants has a decisive effect on brain development. Numerous studies show that dietary fat modulation can change the fatty acid profile of membrane lipid in various neural cells, and therefore influence brain function and development. A diet rich in omega-3 fatty acids increases the level of omega-3 fatty acids in synaptosomal and microsomal membrane lipids in rat brain (20,21). As DHA is essential for the eye and brain function and development in premature infants, feeding premature infants a formula with a high ratio of linoleic acid (LA) to linolenic acid (LNA) (30:1) produced reduced electroretinogram responsiveness early in infancy as compared to those fed human milk or a formula with low ratio of LA to LNA (9:1). Addition of fish oil rich in long-chain omega-3 fatty acids improved some electroretinogram responses as well as visual-acuity development (22). Functional and developmental changes caused by dietary fat may be the result of changes in membrane phospholipid biosynthesis and turnover (21), receptors and transport sites located at membranes and membrane enzymes controlling metabolic processes in brain (23). These changes may also relate to the alteration in ion permeability and nerve transmission (20,21). For example, brain phospholipid fatty acid composition, phosphatidyl ethanolamine methyltransferase acitivity and rate of phosphatidylcholine biosynthesis via the CDP-choline pathway can be manipulated by omega-6 to omega-3 fatty acid ratio. Other studies showed that dietary modulation affects the de novo synthesis of phosphatidylcholine in the developing brain (21,24). Therefore, long-chain omega-3 fatty acids present in preterm human milk

play an important role in the functioning and development of eyes and brain of premature infants, as well as energy production and biosynthesis of essential structural lipids in other body tissues.

The Shortage of Calcium and Phosphorus of Preterm Human Milk

The rapid growth of premature infants may reveal certain nutritional inadequacies of breast milk that are not obvious at slower growth rates. In fact, studies show that although the favorable nitrogen, fat, potassium, sodium, chloride, and energy value of preterm milk may be adequate to meet the nutrient needs of premature infants during the first four weeks of life, premature infants fed exclusively breast milk have shown deficiencies in certain minerals such as calcium, phosphorus, and probably magnesium (25-27). When compared to mature human milk, premature human milk contains relatively less calcium and phosphorus (28). When premature infants are fed unsupplemented breast milk, a 50% reduction in calcium and phosphorus retention has been found compared to a standard commercial formula (29). Since calcium and phosphorus levels in preterm milk decrease significantly over the first four postnatal weeks (1,28,30), a further decrease in the intake of milk calcium after the first postnatal week occurs (1). Therefore, breast milk without supplementation cannot provide adequate and appropriate nutrients for the premature infant (31). Since rapid skeletal growth normally occurs in the last trimester of pregnancy, the lower calcium and phosphorus content of human milk may be insufficient for normal extrauterine mineralization to occur (32-37). In utero, the fetus accumulates 100-200 mg calcium per kilogram body weight

per day by active transport across the placenta between 25 and 36 weeks gestation (34). Therefore approximately 60 mg calcium intake per kilogram body weight per day by premature infants (1) does not meet the calcium requirement needed to achieve intrauterine accretion rates. When calcium accumulation measured for infants fed different milks is compared with the estimated amount of calcium accumulated by the fetus of equivalent gestational age in utero, premature infants fed breast milk have an absolute dietary deficiency of calcium (1,36).

The Evidence of Calcium and Phosphorus Deficiency in Premature infants and Its Improvement after Dietary Supplementations

Calcium is an essential element for body functions. Ninety-nine percent calcium is present in body skeleton, and the other 1% is an important cofactor for neural transmission, enzyme activity, blood coagulation and other extracullular or intracellular functions. Bone mineralization can be affected by calcium and phosphorus metabolism as the endogenous store at birth and ability to absorb and excrete exogenous calcium and phosphorus in infants vary. Calcium and phosphorus deficiency, or rickets, and phosphate depletion syndrome may develop in infants receiving long term breast feeding without supplementation (29,31,38). A lower bone mineralization rate has also been recorded (39). Although calcium absorption increases with increasing postnatal age, intrauterine calcium retention rate is not achieved when infants were fed with breast milk alone (40).

These studies indicate a necessity for early calcium supplementation

in the premature infant fed breast milk (31,33,41). Improvement in bone mineral content and plasma and growth parameters have been observed after supplementation. Disorders such as rickets and phosphate depletion syndrome are readily corrected by calcium and phosphorus supplementation (42). Preterm infants receiving fortified preterm human milk show higher serum calcium and phosphorus status, lower alkaline phosphatase level, and faster weight gain, which may be partially due to better fat absorption of preterm milk as well (43-45). Formula enriched in calcium and phosphorus also enables premature infants to achieve the faster mean weight gain and increment of other growth parameters compared to exclusively pooled preterm milk feeding, which has resulted in a higher alkaline phosphatase levels in these infants (29,46). Bone densitometry studies have shown increased bone density in calcium-supplemented infants (47,48).

3. Factors Influencing Intestinal Fat Absorption

Intraluminal Factors

The bioavailability of bile acids and lipolytic enzymes and the maturity of the digestive system influence fat absorption (2,49-52). Bile acid concentration is inversely correlated with steatorrhea (53) and fecal bile acid excretion is inversely correlated with absorption coefficients of total fat and unsaturated fat (51). Infants with normal absorption coefficients for fatty acids (greater than 80%) usually had a bile acid concentration greater than 4.0 mM, while infants with steatorrhea

(absorption coefficient less than 80%) had a 0-2 mM bile acid concentration (51). Study of fasting duodenal bile acids in premature infants revealed a linear correlation between fat absorption and duodenal bile acid concentration (50,54).

Fat absorption increases with postnatal age (36,55). Lipid absorption in full-term newborn infants is not fully developed (52). This is more apparent in premature infants (56-58) because of the low levels of pancreatic lipases (59,60) and bile salts (59,61). In premature infants, the bile acid pool is reduced to almost 50% of that in full-term infants, which is also correlated with the decreased intraluminal bile acid concentration (61). Infants with lower birthweight usually exibit lower duodenal bile acid concentration compared to older or larger infants (53). Low pancreatic lipase content in full-term newborn infants further limits fat absorption, and this inadequacy is even more obvious in the premature infant (62). Premature infants usually absorb only 65-75% milk fat, which is much lower than the absorption rate of 85-90% of milk fat in full-term infants (63).

However, the presence of preduodenal lipases in neonates, including both lingual lipase and gastric lipase, and bile salt-stimulated lipase present in human milk, partially compensate for the insufficiency of duodenal bile salts and pancreatic lipase in the premature infant, through intragastric lipolysis. Suckling may stimulate secretion of lingual lipase, as nasojejunal feeding causes fat malabsorption in healthy premature infants (64). Nasogastrically fed infants grow faster and show greater head circumferences as compared to nasojejunal feeding, indicating

the significance of gastric lypolysis (65). In rats, progressive increase in intragastric digestion of rat milk triglycerides occurs from birth until weaning. In human subjects, lingual lipase is present before 26 weeks of gestation in the fetus, accumulates in the stomach (66) and is well suited to the acidic condition of stomach (67). Gastric lipase originates in the chief cells of the stomach (68). In premature infants a substantial percentage of triglycerides in milk is rapidly hydrolyzed in the stomach by lingual and gastric lipases. Lingual lipase hydrolyzes medium-chain triglycerides much more efficiently than long-chain triglycerides (69). However, gastric lipase only acts on short chain triglycerides. It has also been found that the sn-3 ester bond is more easily hydrolyzed by lingual lipase than the other two ester sites (70). The lipolytic product is partially comprised of glycerides and free fatty acids, of which the short-chain fatty acids and medium-chain fatty acids can be absorbed directly in the stomach (69). Therefore, intragastric lipolysis contributes significantly to fat absorption. In fact, children with a congenital absence of pancreatic lipase can absorb as much as 60-70% of ingested fat (71).

In premature infants, lingual lipase may remain active in duodenum (56,58,72). Because of the relatively low pH in duodenum of premature infants, considerable lipolytic activity occurs in the duodenum (66) as the enzyme has a wide range of pH stability (pH 4-7) (67). Although the high concentration of bile salts inhibits lingual lipase activity, the lower concentration of bile salts in premature infants does not affect or even stimulate the enzyme activity (66).

Intragastric lipolysis may further facilitate the intestinal hydrolysis of dietary fat by pancreatic lipase (73-76). After incubation with gastric lipase for 30 minutes, triglycerides in human milk are rapidly hydrolyzed by pancreatic enzyme without a lag phase (75). This synergistic effect of two enzymes may be explained as follows. First, the lipolytic products of intragastric digestion, free fatty acids, help to emulsify the remaining triglycerides (73). Second, lingual lipase is more penetrative to milk fat globules and thus the triglyceride core is readily hydrolyzed by the enzyme and forms a better substrate for pancreatic lipase hydrolysis (73,77,78). Third, the free fatty acid produced in intragastric lipolysis may enhance the binding of pancreatic lipase to colipase, and the binding of colipase to bile salts and substrates (79-82). In short, intragastric lipolysis produced by lingual and gastric lipases enhances the affinity of pancreatic lipase and colipase for fat droplets, thus ensuring the efficient digestion of fat in duodenum.

Dietary Factors

Bile salt-stimulated lipase, which is synthesized in the mammary gland of the higher primates (83), plays a significant role in fat absorption of premature infants. It is well suited for this lipolytic process in the infant's intestine as the enzyme is stable in the low pH environment of stomach, and is resistant to pancreatic proteases in the presence of bile salts (84). Heat treatment of human milk inactivates the enzyme and, therefore, reduces fat absorption by about one third in

premature infants (3,58). Vitamin A palmitate is hydrolyzed more rapidly in the infant given fresh human milk when compared to the same infant receiving heat-treated milk (85).

The cooperative function of bile salt-stimulated lipase and pancreatic lipase results in the more complete decomposition of triglycerides in diets. Pancreatic lipase-colipase complex does not hydrolyze dietary triglycerides completely, but instead forms beta-monoglycerides as the main product (86). If the lipolytic product is not removed in time, the lipolysis will slow down and eventually an equilibrium is reached (87). An effect called product inhibition thus results. Addition of bile salt-stimulated lipase rapidly breaks down the monoglyceride ester bond with the formation of free glycerols and fatty acids and thus drives intestinal fat lipolysis toward completion (9). The enzyme is activated by bile salts present in intestinal lumen. The low bile acid concentration in neonatal infants is sufficient to activate the enzyme and thus the fat absorption in breast milk fed premature infants can be much improved in spite of the inadequate concentration of pancreatic lipase and bile salts in these infants (9). Therefore, a strong correlation between fat absorption and intraduodenal bile salt levels has been found in premature infants fed infant formula, but not in infants fed human milk, since the bile salt-stimulated lipase in human milk compensates for the shortage of the endogeneous lipolytic enzyme (50). Thus, infants fed human milk showed more fat absorption compared to those fed formula (50,54).

In addition to the bile salt-stimulated lipase present in human milk,

the characteristics of individual fatty acids and triglycerides in milk contribute to the efficiency of fat absorption. The position of a fatty acid on the glycerol moiety in a triglyceride can limit fat absorption (2,51,88). Most palmitate in fresh human milk is esterified in the beta-position of glycerol molecules and remains unhydrolyzed by lipase. The beta-monoglyceride thus formed is absorbed directly instead of forming an insoluble palmitic acid-calcium soap in intestine (2). In rats, a linear correlation between fat absorption and the proportion of a fatty acid in the beta-position is observed for palmitic, and to a lesser extent, for myristic and oleic acid (89). When newborn infants were given similar formulas in which palmitic acid was either randomized on the triglyceride molecule or located mainly at the beta-position, fat absorption was markedly reduced when palmitic acid was distributed at random, indicating triglyceride configuration is an important factor in improving fat absorption (90).

The chain length and degree of saturation of individual fatty acids also determine the rate of fat absorption. Unsaturated fatty acids have higher absorption coefficients (51). The absorption of saturated fatty acids varies inversely with chain length (51,91) and medium-chain triglycerides are better absorbed than long-chain triglycerides (51,55,92). It has been estimated that an approximately 10% decrease in fatty acid absorption occurs for each two carbon addition in fatty acid chain length (51), as aqueous solubility of a fatty acid decreases with increasing chain length. In other words, because medium-chain saturated fatty acids are relatively water soluble, they may not require micellar

solubilization by bile acids. Therefore, its absorption coefficient shows no linear correlation with bile acid concentration, or it is efficiently absorbed irrespective of the intraduodenal bile acid concentration (51). For instance, premature infants fed a medium-chain triglyceride formula have a markedly decreased stool volume and frequency and increased fat absorption rate (55,91-93). Long-chain triglycerides are rehydrolyzed prior to absorption and have a relatively poor solubility (2). Thus, fat absorption is very dependent on the fatty acid profile of feedings (88,94-96). Compared to formula containing long-chain triglycerides, an isocaloric formula containing medium-chain triglycerides significantly promotes weight gain and nitrogen retention, in addition to improving fat absorption rate (97).

As discussed above, total fat absorption and calcium absorption are closely correlated (41,49,88,92,95). The mean fat excretion value in infants receiving calcium supplementation is compared with the value in infants not receiving calcium, and it has been found that the fat excretion is increased with calcium supplementation (1,41,48,49,94). Long-chain saturated fatty acids form calcium-soap and increase both fat and calcium excretion (52), or excessive calcium interferes with fat absorption and individual fatty acids selectively (48,94), as palmitic and stearic acids have been identified previously as the major fecal fatty acids in the adult while oleic acid is preferentially absorbed (98).

4. Factors Influencing Intestinal Calcium Absorption

Dietary Calcium and Phosphorus Intake and Their Ratio

The absolute amount of calcium and phosphorus intake and their ratio play a significant role in calcium secretion and absorption (36,41,48,99-103). In suckling rats, net transport of calcium is related in a linear manner to the concentration of calcium in the intestinal lumen (6,104,105). Calcium secretion occurs when intraluminal concentration of calcium is lower than plasma ionized calcium concentration (104). Phosphorus supplementation with calcium, in order to avoid hypercalciuria and phosphate depletion syndrome, improves calcium retention (41,106). Since some of the retained phosphorus is used for new body cell synthesis in premature infants, less phosphorus is available for bone mineralization. Calcium is thus accumulated in the plasma of these breast milk fed infants and results in hypercalcemia and hypercalciuria, which can be readily corrected by phosphorus supplementation (107). A positive correlation between serum calcium concentration and urinary calcium excretion and a negative correlation between serum calcium concentration and serum phosphorus concentration or urinary phosphorus excretion have also been observed (107). Therefore, a Ca/P ratio of 1.7-2.0:1 has been suggested to approximately achieve the intrauterine bone mineralization curve in premature infants (108).

Vitamin D and Other Regulating Factors

Homeostatic regulation of calcium depends on vitamin D, parathyroid

hormone, and calcitonin. These hormones act on the intestine, kidney, and bone to maintain serum calcium at a physiological level. Vitamin D, 1,25-$(OH)_2$ D_3 controls calcium metabolism through calcium intestinal absorption, bone mobilization and renal reabsorption. It increases serum calcium and phosphorus level for normal bone mineralization by improving intestinal calcium absorption, bone reabsorption and renal reabsorption at the distal tubule.

The saturable calcium transport, predominant in proximal intestine, is regulated by vitamin D. This saturable process consists of entry of calcium across the brush-border membrane, intracellular diffusion, and extrusion across the basolateral membrane (109). Calcium transport across cell membrane is carrier-mediated and inhibited competitively by strontium and noncompetitively by magnesium (110). As membrane lipid composition determines the function of a mobile carrier in a lipid bilayer, liposome treated brush-border membrane vesicles showed a change in phospholipid and cholesterol and, consequently, a decrease in Vmax which is correlated with brush-border membrane vesicle cholesterol content and fluidity (110). Vitamin D can affect phospholipid metabolism of intestine, kidney, bone, and possibly parathyroid cells (111). Vitamin D also stimulates active calcium transport by synthesis of a cytosolic protein, calcium-binding protein, and acceleration of intracellular diffusion of calcium. In adults, 1,25-$(OH)_2$ D_3 stimulates this energy-dependent cellular process in intestinal epithelium and converts regions of net calcium secretion in ileum and colon to net absorption (112). In weanling and post-weanling rats, vitamin D enhances active calcium uptake, significantly elevating

plasma calcium (6).

Vitamin D is activated in the liver and kidney. The transformation is stimulated by decreases in serum calcium and phosphorus and increase in serum parathyroid hormone. The active form of vitamin D, 1,25 $(OH)_2$ D_3, then binds to a receptor in the nuclear membrane. The complex further binds to specific portions of nuclear DNA, causing transcription of specific genes that code for calcium and phosphorus transport proteins. Calcium-binding protein has been identified in the cytoplasm, which facilitates calcium diffusion from the microvillar pole of the enterocyte to the basolateral membrane. The rate of intracellular calcium diffusion in the absence of calcium-binding protein is only about 1/70 of that found in the vitamin D-replete cell (109). In vitamin D-deficient or vitamin D-replete chicks, giving vitamin D at normal calcium and phosphorus levels results in an elevation of duodenal calcium-binding protein and its mRNA levels (113,114). The low level of duodenal vitamin D-dependent calcium-binding protein in hypophosphatemic mice is associated with low calcium intestinal absorption and low bone mineralization (115). Vitamin D deficiency can also cause reduction in body weight, intestinal mucosa weight, total protein and DNA and some hydrolytic enzyme activities both in intestine and pancreas (116).

In premature human neonates, calcium absorption and retention may be increased by adding vitamin D, especially after the first week of life (42,106,117,118). It seems that intestinal calcium absorption is not very responsive to vitamin D feeding in the early days of life although vitamin D can be well absorbed in premature infants (117). Indeed, premature

infants with hypocalcemia did not show vitamin D activation in the first week of life (119). It is hypothesized that in the early days of life, a nonsaturable, or passive calcium transport process is predominant, which may be vitamin D-independent (105). Therefore, the calcium status at this stage is maintained primarily by the relatively high calcium content present in milk, endocytosis of ingested calcium or simple passive diffusion down the calcium concentration gradient. The presence of lactose in milk increases passive absorption of calcium in intestine (6). It might be possible that the intestine is not sensitive to vitamin D in the early days of life because of the lack of vitamin D receptor in the nuclear membrane.

Parathyroid hormone is another regulating factor. The effect of vitamin D on bone mobilization and renal reabsorption of calcium requires the presence of this hormone. Its synthesis and secretion are stimulated by the reduction in serum calcium and 1,25 $(OH)_2D_3$ levels. Parathyoid hormone acts directly on bone and kidney to increase extracellular calcium concentration and indirectly on intestine to increase calcium absorption by stimulating vitamin D activation in kidney (120). As the secretion of parathyroid hormone may be mediated by the magnesium concentration in this manner, magnesium might affect calcium metabolism as well.

The elevation in serum calcium stimulates the production and release of calcitonin from the parafollicular cell of thyroid gland. Calcitonin decreases serum calcium level by reducing bone resorption and tubular reabsorption of calcium and inactivation of vitamin D by modulating some enzyme activities such as adenyl cyclase, and vitamin D 1-alpha-

hydroxylase. Calcitonin also indirectly inhibits calcium absorption in intestine by decreasing gastrin and gastric acid secretion (120).

Dietary Fat

Calcium absorption is also related to fat absorption. As calcium salts are formed with fat and excreted in increased amounts when calcium is supplemented to feeds (1,2,41,48,94,117), intestinal calcium absorption is favored by complete fat absorption by pancreatic enzymes. Long-chain saturated fatty acids are mainly excreted in the form of insoluble soaps. Infants fed a formula containing medium-chain triglycerides absorb more calcium than infants fed a standard cow's milk-based formula (95). The enhanced calcium absorption is proportional to the medium-chain triglyceride concentration in the formula and is positively correlated with fat absorption (95). In other words, in newborn infants, fat excreted as calcium-soap is related to fatty acid composition of a formula (121). A high level of stearate and palmitate in formula increases calcium losses, and retention of calcium is greatest in a formula low in stearate and palmitate and high in oleate (121). In chicks fed diets supplemented with palmitic acid, a significant reduction in bone ash and bone calcium content has been found (122). Increasing the calcium in the diet further decreases calcium retention and bone calcium content associated with addition of fat (122).

Other Factors Affecting Calcium Absorption

Some electrolytes affect calcium transport and absorption in the

intestinal lumen or mucosa and renal glomeruli. Sodium induces hypercalciuria and alters calcium absorption in intestine and reabsorption in kidney (112,123,124). In intestine, sodium affects calcium absorption at the brush border by altering the transmembrane electrical gradient and at the basolateral membrane by exchanging with intracellular calcium (112). Sodium intake in the hypocalcemic premature infant is positively correlated with urinary calcium, phosphorus, and magnesium excretions (124). Moreover, magnesium inhibits calcium absorption by depressing passive calcium absorption in the jejunum of adult human subjects (125).

Some drugs have also been shown to influence calcium metabolism. Furosemide induces hypercalciuria (123,126,127), and bone demineralization and renal calcification are found in premature infants receiving this diuretic treatment (126,127). Glucocorticoids suppress calcium absorption in intestine in the suckling rat (128). Calcium absorption also increases with gestational age and postnatal age (36,55,102). As vitamin D can be digested by bile salt, inadequacy of luminal bile salt may decrease calcium absorption as well. Acidity of digestate may enhance calcium absorption, therefore, some sugars increase intestinal calcium absorption by lowering luminal pH and thus increasing the solubility of calcium (129).

5. Introduction to Phosphorus Absorption

The low phosphorus content of human milk, especially preterm human milk, and gradual decline of phosphorus over the first few weeks of lactation are insufficient for growing low birthweight infants (28,29,31).

Premature infants excrete more phosphorus in urine in the first week of life compared to term infants (130). These infants exhibit decreased serum phosphorus and increased alkaline phosphase levels (29). Phosphorus or phosphorus and calcium supplementation can correct this situation, as phosphorus retention, bone content, and weight gain have been improved (41,108,131-133). As phosphate depletion is related to neonatal hypercalciuria in the premature infant fed breast milk (107), it is necessary to briefly review the intestinal phosphorus absorption and its interactions with calcium and magnesium.

Eighty five per cent of phosphorus deposits are found in skeleton tissue, the other 15% are present in soft tissues (120). Its metabolism is mainly regulated by vitamin D and parathyroid hormone (120). Parathyroid hormone decreases phosphorus reabsorption in kidney. Vitamin D increases renal tubular reabsorption and intestinal absorption of phosphorus. Tubular reabsorption and serum concentration of phosphorus can be also decreased by calcitonin.

Several factors may affect phosphorus absorption and retention. Increasing amount of phosphorus in milk may increase phosphorus retention (102). Calcium supplementation increases not only calcium absorption but also causes a dose-dependent phosphorus retention in premature infants (41,102,134). Magnesium in diets may decrease phosphorus absorption by forming insoluble salt with magnesium in intestinal lumen. Phosphorus uptake by the intestinal mucosa decreases with aging. Although vitamin D and calcium intake can stimulate phosphorus uptake (135), the capacity is most intensified in young animals (136). Phosphorus absorption and

retention is also enhanced by a medium-chain fatty acid-enriched diet (50%) in adult rats receiving intestinal resection (137). Aluminum commonly contained in antacids interferes with intestinal phosphorus absorption, resulting in phosphate depletion (138). Urinary phosphorus excretion can be increased by ingestion of anions irrespective of their sodium or potassium salts (139).

6. Introduction to Magnesium Absorption

When compared to formulas, preterm human milk has a relatively low concentration of magnesium, which does not enable premature infants to achieve intrauterine magnesium retention rate (31).

More than half of total body magnesium is present in skeleton, whereas the rest is found in soft tissues. Magnesium is a very important constituent in bone mass. It is also involved in various enzymatic reactions in both energy metabolism and synthesis of proteins and nuclear acids.

Magnesium is absorbed throughout the intestine, mainly distal small intestine. The mechanism of the intestinal absorption includes passive diffusion, solvent drag, and active transport, with the passive diffusion being the major pathway and, possibly, a limited active transport in descending colon in some adult or aged animals (140). In fact, in suckling or adolescent rats, no active transport of magnesium is found. Extracellular magnesium is filtered at the glomerulus and most filtered magnesium is reabsorbed in the proximal tubule and thick ascending limb.

Various factors can influence magnesium absorption or retention.

There are also complex interactions between the divalent cations in intestinal lumen and complicated effect of vitamin D on magnesium (140). Increased magnesium intake is associated with increased absorption and bone accumulation (141). Duodenal magnesium absorption is suggested to be totally concentration-dependent (142). Pharmacological doses of vitamin D administration increases magnesium absorption and serum magnesium concentration (140,143), whereas a substantial amount of magnesium absorption is independent of vitamin D (140). As vitamin D also increases urinary magnesium excretion, the retention rate of magnesium may not be greatly affected by vitamin D (140). In patients with hypomagnesemia after intestinal resection, magnesium status is improved by vitamin D and magnesium supplementation (144).

Dietary calcium interferes with magnesium absorption, possibly by competition for a common carrier system, modulation of membrane permeability to magnesium or a specific magnesium carrier, direct competition in passive transport, or formation of a complex agent with each other. It is also possible that this interaction, to a certain extent, is conducted by hormonal feedback, as both vitamin D and parathyroid hormone are very responsive to the serum concentration of calcium and magnesium. Phosphorus in the diet also affects magnesium absorption by forming insoluble salt with magnesium, resulting in greater magnesium excretion in stool. In one study with premature infants fed formula with calcium and phosphorus supplementation, magnesium absorption and retention were severely impaired and showed a negative balance (145).

Some other dietary factors affect intestinal magnesium absorption.

Phytate in soybean products decreases magnesium absorption and bone content compared to casein-predominant formulas because of Mg-Ca-phytate complex formation in the intestinal lumen (141). Lactose present in milk decreases luminal pH through intestinal microbial fermentation and thus enhances magnesium solubility and passive magnesium absorption (141). Intraluminal potassium depresses magnesium absorption (146,147). A diet supplemented with anions increases urinary magnesium excretion and, therefore, the retention rate is reduced (139). In rats, short-chain fatty acids derived from fermentation of carbohydrate in the large intestine stimulate magnesium absorption by offering protons to Mg^{++}/H^{+} in the apical membrane of epithelium (147). Magnesium secretion in proximal intestine is greatly increased in rats with intestinal resection (148).

Magnesium deficiency in the diet causes hypocalcemia and hypomagnesemia. Vitamin D administration improves both calcium and magnesium status (143). It is suggested that magnesium deficiency impairs synthesis or target organ response to vitamin D (143). A protective effect of magnesium in renal calcinosis in furosemide-treated rats with magnesium deficiency has also been suggested (149).

7. Summary

Preterm human milk is recommended for the premature infant as it contains more medium-chain fatty acids, long-chain polyunsaturated fatty acids, and bile salt-stimulated lipase, therefore ensuring superior fat absorption. Long-chain omega-3 fatty acids in human milk are essential to the function and development of premature infants' eyes and brain.

However, the calcium and phosphorus content of breast milk is insufficient for normal bone mineralization in the premature infant. Therefore, calcium supplementation is necessary to obtain an intrauterine calcium accretion rate. Calcium supplementation increases lipid loss in stool as it forms calcium soap with long-chain saturated fatty acids in intestinal lumen. Factors influencing calcium absorption include absolute intake of calcium and phosphorus and their ratio, vitamin D status, dietary fat, some cations and drugs, and gestational and postnatal age. Fat absorption can be affected by the presence and concentration of bile salts, lipolytic enzymes, postnatal age, calcium intake, properties of milk lipids such as fatty acid saturation, chain length, and the position of fatty acids in glycerol molecules. Poor fat absorption in the premature infant due to lower intestinal concentration of bile salts and pancreatic lipase can be improved with breast milk feeding because of the compensation of intragastric lipolysis by lingual lipase and gastric lipase and intestinal hydrolysis by bile salt-stimulated lipase present in human milk. Lingual lipase does not only hydrolyze fat in stomach but also aids intestinal fat absorption. As well, bile salt-stimulated lipase makes intestinal lipolysis more complete. Intestinal phosphorus absorption may be enhanced by dietary intake of phosphorus, calcium, and medium-chain fatty acids and diminished by dietary magnesium, aluminum and some anions. Intestinal magnesium absorption can be improved by magnesium, short-chain fatty acids, and lactose in diets and impaired by dietary calcium, phosphorus, phytate, potassium and some anions. Vitamin D, parathyroid hormone and calcitonin regulate calcium, phosphorus and magnesium metabolism by acting

on the intestine, kidney and bone and thus maintain their serum concentrations at physiological levels.

C: Rationale

Because of the superior nutritional and immunological value of human milk, feeding premature infants' their own mother's milk has been recommended, especially in the first few weeks of life. However, calcium and phosphorus deficiency is quite evident in the premature infant fed exclusively human milk, as the calcium and phosphorus content in human milk is relatively low and mineral accumulation of the premature infant is very limited. Therefore, calcium or calcium and phosphorus supplementation is essential for the premature infant to reach the intrauterine mineral accretion rate. Unfortunately, calcium soap formation with fatty acids, especially long-chain saturated fatty acids complicates this practice, as the energy intake and, possibly, some essential fatty acids and consequently growth and development of the premature infant will be affected by the increased excretion of calcium-fatty acid soaps following calcium supplementation.

It is obvious that a more appropriate method of calcium supplementation requires investigation. In most related studies, calcium supplements have been given orally with feedings of either human milk or formula. In this study, an alternative way of calcium supplementation was compared with this practice, i.e. giving the supplement between two feedings to the premature infant fed expressed preterm human milk. The study will, to a certain extent, further clarify the interactions of

lipids and calcium and other divalent cations which may interfere with calcium metabolism.

1. Hypothesis

It is hypothesized that by changing the timing of calcium supplementation from with feedings to between feedings in the premature infant fed breast milk, insoluble calcium salt formation will be decreased. It is hypothesized that both fat and calcium losses in stool will decrease and the absorption coefficient of long chain saturated fatty acids will increase in the infant receiving calcium supplementation between feedings.

2. Objectives

a. To investigate the effect of timing of calcium supplementation on calcium and fat absorption, and absorption coefficients of selected fatty acids in the premature infant fed breast milk.

b. To investigate the interaction of intestinal minerals (Ca, P, Mg) and fatty acids in orally fed premature infants.

E: References

1. Atkinson S, Clandinin MT. Calcium supplementation of mother's milk for low birthweight infants: problems related to absorption and retention. Nutrition Res. 7: 813-23, 1987.

2. Alemi B, Hamosh M, Scanlon JW, Salxman-Mann C, Hamosh P. Fat digestion in VLBW infants: effect of addition of human milk to low-birth-weight formula. Pediatrics 68: 484-9, 1981.

3. Atkinson S, Bryan M, Anderson GH. Human milk feeding in premature infants: protein, fat and carbohydrate balances in the first two weeks of life. J Pediatr. 99: 617-24, 1981.

4. Chessex P, Reichman B, Verellen G, Putet G, Smith JM, Hein T, Swyer P. Quality of growth in premature infants fed their own mother's milk. J Pediatr. 102: 107-12, 1983.

5. Lemons JA, Moye L, Hall D, Simmons M. Differences in the composition of preterm and term human milk during early lactation. Pediatr Res. 16: 113-7, 1982.

6. Halloran B, Deluca H. Calcium transport in small intestine during early development: role of vitamin D. Am J Physiol. 239: G473-9, 1980.

7. Hamosh M. Lipid metabolism in premature infants. Biol Neonate 52 Suppl. 1: 50-64, 1987.

8. Pettifor JM, Stein H, Herman A, Ross FP, Blumenfeld T, Moodley GP. Mineral homeostasis in very low birth weight infants fed either own mother's milk or pooled pasteurized preterm milk. J Pediatr Gastroenterol Nutr. 5: 248-53, 1986.

9. Hernell O, Blackberg L. Digestion of human milk lipids: physiologic significance of sn-2 monoacylglycerol hydrolysis by bile salt-stimulated lipase. Pediatr Res. 16: 882-5, 1982.

10. Gross SJ. Growth and biochemical response of preterm infants fed human milk or modified infant formula. N Engl J Med. 3; 308 (5): 237-41, 1983.

11. Simopoulos AP. Omega-3 fatty acids in growth and development and in health and disease. Part I: the role of omega-3 fatty acids in growth and development. Nutr Today 23 (2): 10-9, 1988.

12. Clandinin MT. Feeding and absorption of long-chain fatty acids in very-low-birth-weight infants. Nutritional Requirements of the Low-

Birthweight Neonate. Excerpta Medica pp.41-50, 1987.

13. Clandinin MT, Chappell JE, Van Aerde J. Requirements of newborn infants for long-chain polyunsaturated fatty acids. Acta Pediatr Scand Suppl. 351:63-71, 1989.

14. Martinez M. Dietary polyunsaturated fatty acids in relation to neural development in humans. In: Galli C, Simopoulos AP. Dietary omega-3 and omega-6 fatty acids: biological effects and nutritional essentiallity. New York: Plenum Press. pp.123-33, 1989.

15. Walker BL. Maternal diet and brain fatty acids in young rats. Lipids 2: 497-500, 1967.

16. Connor WE, Neuringer M, Reisbick S. Essentiality of omega-3 fatty acids: evidence from the primate model and implications for human nutrition. In: Simopoulos AP, Kifer RR, Martin RE, Barlow SM. Health effects of omega-3 polyunsaturated fatty acids in seafoods. World Rev Nutr Diet. 66: 118-32, 1991.

17. Bazan NG. The supply of omega-3 polyunsaturated fatty acids to photoreceptors and synapses. In: Galli C, Simopoulos AP. Dietary omega-3 and omega-6 fatty acids: biological effects and nutritional essentiality. New York: Plenum Press. pp.227-40, 1989.

18. Bourre JM, Dumont O, Piciotti M, Pascal G, Durand G. Polyunsaturated fatty acids of the n-3 series and nervous system development. In: Galli C, Simopoulos AP. Dietary omega-3 and omega-6 fatty acids: biological effects and nutritional essentiality. New York: Plenum Press. pp.159-76, 1989.

19. Rao PS. Fatty acid composition of cerebrosides and phospholipids in brains of undernourished rats. Nutr Metab. 23: 136, 1979.

20. Foot M, Cruz TF, Clandinin MT. Influence of dietary fat on the lipid composition of rat brain synaptosomal and microsomal membranes. Biochem J. 208: 631-40, 1982.

21. Hargreaves K, Clandinin MT. Phosphatidylethanolamine methyl-transferase: evidence for influence of diet fat on selectivity of substrate for methylation in rat brain synaptic plasma membranes. Biochim Biophys Acta 918: 97-105, 1987.

22. Uauy RD, Birch DG, Birch EE, Tyson JE, Hoffman DR. Effect of dietary omega-3 fatty acids on retinal function of very-low-birth-weight neonates. Pediatr Res. 28: 485-92, 1990.

23. Hargreaves K, Clandinin MT. Phosphocholinetransferase activity in plasma membrane: effect of diet. Biochem Biophys Res Comm. 145:

309-15, 1987.

24. Hargreaves K, Clandinin MT. Co-ordinate control of CDP-choline and phosphatidylethanolamine methyltransferase pathways for phosphatidylcholine biosynthesis occurs in response to change in diet fat. Biochim Biophys Acta 1001: 262-7, 1989.

25. Atkinson SA, Anderson GH, Bryan MH. Human milk: Comparison of the nitrogen composition in milk from mothers of premature and full-term infants. Am J Clin Nutr. 33: 811-5, 1980.

26. Atkinson SA, Radde IC, Chance GW, Bryan MH, Anderson GH. Macromineral content of milk obtained during early lactation from mothers of premature infants. Early Hum Dev. 4: 5-14, 1980.

27. Gross SJ, David RJ, Bauman L, Tomarelli RM. Nutritional composition of milk produced by mothers delivering preterm. J Pediatr. 96: 641-4, 1980.

28. Butte NF, Garza C, Johnson CA, Smith EO'B, Nichols BL. Longitudinal changes in milk composition of mothers delivering preterm and term infants. Early Hum Dev. 9: 153-62, 1984.

29. Jowe J, Rowe D, Horak E, Spackman T, Saltzman R, Robinson S, Philipps A, Raye J. Hypophosphatemia and hypercalciuria in small premature infants fed human milk: evidence for inadequate dietary phosphorus. J Pediatr. 104 (1): 112-7, 1984.

30. Chan GM. Human milk calcium and phosphate levels of mothers delivering term and preterm infants. J Pediatr Gastroenterol Nutr. 1 (2): 201-5, 1982.

31. Atkinson SA, Radde IC, Anderson GH. Macromineral balances in premature infants fed their own mothers' milk or formula. J Pediatr. 102 (1): 99-106, 1983.

32. Rowe J, Wood D, Eowe D, Raisz L. Nutritional hypophosphatemic rickets in a premature infant fed breast milk. N Eng J Med. 300: 293, 1979.

33. Callenbach H, Sheehan M, Abramson S, Hall R. Etiologic factors in rickets of very low birth weight infants. J Pediatr. 98: 800, 1981.

34. Widdowson Elsie M. Changes in body composition during growth. Scientific Foundation of Pediatrics 575: 332-3, 1981.

35. Fleischman A, Perk H, Skolnick L. Rickets in the small premature. Pediatr Res. 14: 571, 1980.

36. Shaw JCL. Evidence for defective skeletal mineralization in LBW infants: the absorption of calcium and fat. Pediatrics 57: 16-25, 1976.

37. Chan GM, Mileur LJ. Posthospitalization growth and bone mineral status of normal preterm infants. Feeding with mother's milk or standard formula. Am J Dis Child. 139 (9): 896-8, 1985.

38. Greer FR, Steichen JJ, Tsang RC. Calcium and phosphate supplements in breast milk related rickets. Am J Dis Child. 136: 581-3, 1982.

39. Chan GM, Mileur L, Hansen JW. Effects of increased calcium and phosphorous formulas and human milk on bone mineralization in preterm infants. J Pediatr Gastroenterol Nutr. 5: 444-9, 1986.

40. Schanler RJ, Abrams SA, Garza C. Mineral balance studies in very low birthweight infants fed human milk. J Pediatr. 113: 230-8, 1988.

41. Salle B, Senterre B, Putet G, Rigo J. Effects of calcium and phosphorus supplementation on calcium retention and fat absorption in preterm infant fed pooled human milk. J Pediatr Gastroenterol Nutr. 5: 638-42, 1986.

42. Senterre J, Putet S, Salle B, Rigo J. Effects of vitamin D and phosphorus supplementation on calcium retention in preterm infants fed pooled human milk. J Pediatr. 103: 305-7, 1983.

43. Modanlou HD, Lim MO, Hansen JW, Sickles V. Growth, biochemical status, and mineral metabolism in very-low-birth-weight infants receiving fortified preterm human milk. J Pediatr Gastroenterol Nutr. 5: 762-7, 1986.

44. Gross SJ. Bone mineralization in preterm infants fed human milk with and without mineral supplementation. J Pediatr. 111: 450-8, 1987.

45. Greer FR, Mc Cormick A. Improved bone mineralization and growth in premature infants fed fortified own mother's milk. J Pediatr. 112: 961-9, 1988.

46. Cooper PA, Rothberg AD, Pettifer JM, Bolton KD, Devenhuis S. Growth and biochemical response of premature infants fed pooled preterm milk or special formula. J Pediatr Gastroenterol Nutr. 3(5): 749-54, 1984.

47. Rowe J, Rowe D, Horak E, Sparkman T, Saltzman R, Robinson S, Phillipps A, Raye J. Hypophosphatemia and hypercalciuria in small premature infants fed human milk: evidence of inadequate dietary phosphorus. J Pediatr. 104: 112-7, 1984.

48. Day G, Chance G, Radde I, Park E, Sheepers J. Growth and mineral metabolism in very low birth weight infants. II. Effects of calcium supplementation on growth and divalent cations. Pediatr Res. 9: 568-75, 1975.

49. Katz L, Hamilton JR. Fat absorption in infants of birthweight less than 1300g. J Pediatr. 85: 608-14, 1974.

50. Singer E, Murphy GM, Edkins S, Anderson CM. Role of bile salts in fat absorption of premature infants. Arch Dis Child. 49: 174-80, 1974.

51. Finley AJ, Davidson M. Bile acid excretion and patterns of fatty acid absorption in formula-fed premature infants. Pediatrics 65: 132-8, 1980.

52. Watkins JB, Bliss CM, Donaldson RM, Lester R. Characterization of newborn fecal lipid. Pediatrics 53: 511-5, 1974.

53. Lavy U, Silverberg M, Davidson M. Role of bile acids in fat absorpton in low birth weight infants. Pediatr Res. 5: 387A (abstr.), 1971.

54. Jarvenpaa AL. Feeding the low-birthweight infant. IV. fat absorption as a function of diet and duodenal bile acids. Pediatrics 72: 684-98, 1983.

55. Okamoto E, Muttart C, Zucker C, Heird W. The use of medium chain triglycerides in feeding the low-birth-weight infant. Am J Dis Child. 136: 428-31, 1982.

56. Davidson M, Bauer CH. Patterns of fat excretion in feces of premature infants fed various preparations of milk. Pediatrics 25: 375-84, 1960.

57. Watkins JB. Mechanisms of fat absorption and the development of gastrointestinal function. Pediatr Clin Nutr Am. 22: 721-30, 1975.

58. Williamson S, Finucane E, Ellis H, Gamsu HR. Effect of heat treatment of human milk on absorption of nitrogen, fat, sodium, calcium and phosphorus by preterm infants. Arch Dis Child. 53: 555-63, 1978.

59. Norman A, Strandvik B, Ojamae O. Bile acids and pancreatic enzymes during absorption in the newborn. Acta Pediatr Scand. 61: 571-6, 1972.

60. Fredrikzon B, Olivecrona T. Decrease of lipase and esterase activities in intestinal contents of newborn infants during test

meals. Pediatr Res. 12: 631-4, 1978.

61. Watkins JB, Sczcepanik P, Gould JB, Klein P, Lester R. Bile salts kinetics in premature infants: an explanation for inefficient lipid absorption. Gastroenterology 69: 706-13, 1975.

62. Rey J, Schmitz J, Amedee-Manasme O. Fat absorption in low birthweight infants. Acta Paediatr Scand Suppl. 296: 81-4, 1982.

63. Fomon SJ, Ziegler EE, Thomas LN, et al. Excretion of fat by normal full-term infants fed various milks and formulas. Am J Clin Nutr. 28: 1299-313, 1970.

64. Roy RN, Pollnitz RP, Hamilton JR, Chance GW. Impaired assimilation of nasojejunal feeds in healthy low-birthweight newborn infants. J Pediatr. 90: 431-4, 1977.

65. Whitfield MF. Poor weight gain of the low birthweight infant fed nasojejunally. Arch Dis Child. 57: 597-601, 1982.

66. Hamosh M, Scanlon JW, Ganot D, Likel M, Scanlon KB, Hamosh P. Fat digestion in the newborn: characterization of lipase in gastric aspirates of premature and term infants. J Clin Invest. 67: 838-46, 1981.

67. Cohen M, Morgan GRH, Hofmann AF. Lipolytic activity of human gastric and duodenal juice against medium and long-chain triglycerides. Gastroenterology 60: 1-15, 1971.

68. DeNigris SJ, Hamosh M, Kasbekar DK, Lee TC, Hamosh P. Lingual and gastric lipases: species differences in the origin of prepancreatic digestive lipases and in the localization of gastric lipase. Biochim Biophys Acta 959: 38-45, 1988.

69. Liao Th, Hamosh M, Scanlon JW, Hamosh P. Preduodenal fat digestion in the human infant: effect of fatty acid chain length on triglyceride hydrolysis. Clin Res. 28: 820A, 1980.

70. Jensen RG, De Jong FA, Clark RM, Palmgren LG, Liao TH, Hamosh M. Stereospecificity of premature human infant lingual lipase. Lipids 17: 570-2, 1982.

71. Muller DPR, McCollum JPK, Trompeter RS, Harries JT. Studies on the mechanism of fat absorption in congenital isolated lipase deficiency. Gut 16: 138, 1975.

72. Zoppi G, Andreotti G, Pajno-Derrara F, Njai DM, Gaburro D. Exocrine pancreas function in premature and full-term neonates. Pediatr Res. 6: 880-6, 1972.

73. Plucinski TM, Hamosh M, Hamosh P. Fat digestion in the rat: role of lingual lipase. Am J Physiol. 237: E541-7, 1979.

74. Olivecrona T, Billstrom A, Fredrikzon B, Johnson O, Samuelson G. Gastric lipolysis of human milk lipids in infants with pyloric stenosis. Acta Pediatr Scand. 62: 520-2, 1973.

75. Bernback S, Blackberg L, Hernell O. Fatty acids generated by gastric lipase promote human milk triacylglycerol digestion by pancreatic colipase-dependent lipase. Biochim Biophys Acta 1001: 286-93, 1989.

76. Gargouri Y, Pieroni G, Riviere C, Lowe PA, Sauniere JF, Sarda L, Verger R. Importance of human gastric lipase for intestinal lipolysis: an in vitro study. Biochim Biophys Acta 879: 419-23, 1986.

77. Hamosh M, Scow RO. Lingual lipase and its role in the digestion of dietary fat. J Clin Invest. 52: 88-95, 1973.

78. Edwards-Webb JD, Thompson SY. Studies on lipid digestion in the preruminant calf. 3. The action of salivary lipase on milk fat in the abomasum. Br J Nutr. 40: 125-31, 1978.

79. Patton JS, Donner J, Borgstrom B. Lipase-colipase interactions during gel filtration high and low affinity binding situations. Biochim Biophys Acta 529: 67-78, 1978.

80. Suave P, Desnuelle P. Interactions of pancreatic colipase with taurodeoxycholate-oleate mixtures above the critical micell concentration. FEBS Lett. 122: 91-4, 1980.

81. Borgstrom B. The importance of phospholipids, pancreatic phospholipase A2, fatty acid and other factors for the digestion of dietary fat: in vitro experiments with porcine enzymes. Gastroenterology 78: 954-62, 1980.

82. Larsson A, Erlanson-Albertsson C. The identity and properties of two forms of activated colipase from porcine pancrease. Biochim Biophys Acta 664: 538-48, 1981.

83. Hamosh M. Lingual and breast milk lipases. Adv Pediatr. 29: 33-67, 1982.

84. Hernell O. Human milk lipases. III. Physiological implication of the bile salt-stimulated lipase. Eur J Clin Invest. 5: 267-72, 1975.

85. Fredrikzon B, Hernell O, Blackberg L, Olivecrona T. The bile salt-stimulated lipase in human milk. Evidence of activity in vivo and of

a role in the digestion of milk retinol esters. Pediatr Res. 16: 1048-52, 1982.

86. Borgstrom B. On the mechanism of pancreatic lipolysis of glycerides. Biochim Biophys Acta 13: 491-504, 1954.

87. Borgstrom B. Influence of bile salt, pH, and time on the action of pancreatic lipase: physiological implications. J Lipid Res. 5: 522-31, 1964.

88. Barnes L, Morrow G, Silverio J, Finnegan L, Heitman S. Calcium and fat absorption from infant formulas with different fat blends. Pediatrics 54: 217-21, 1974.

89. Tomarelli RM, Meyer BJ, Weaber JR, Bernhart FW. Effect of positional distribution on the absorption of the fatty acids of human milk and infant formulas. J Nutr. 95: 583-90, 1968.

90. Filer LS, Mattson FH, Fomon SJ. Triglyceride configuration and fat absorption by the human infant. J Nutr. 99: 293-8, 1969.

91. Jensen C, Buist N, Wilson T. Absorption of individual fatty acids from long-chain or medium-chain triglycerides in very small infants. Am J Clin Nutr. 43: 745-51, 1986.

92. Huston R, Reynolds J, Jensen C, Buist N. Nutrient and mineral retention and vitamin D absorption in low-birth-weight infants: effect of medium-chain triglycerides. Pediatrics 72: 44-8, 1983.

93. Tantibhedhyangkul P, Hashim SA. Medium-chain triglyceride feeding in premature infants: effects on fat and nitrogen absorption. Pediatrics 55: 359-70, 1975.

94. Chappell J, Clandinin MT, Kearney-Volpe C, Reichman V, Swyer P. Fatty acid balance studies in premature infants fed human milk or formula: effect of calcium supplementation. J Pediatr. 108: 439-47, 1986.

95. Tantibhedhyangkul P, Hashim S. Medium-chain triglyceride feeding in premature infants: effect on calcium and magnesium absorption. Pediatrics 61: 537-45, 1978.

96. Van Aerde J, Sauer P, Heim T, Smith J, Swyer P. The effect of diet composition on energy metabolism, substrate utilization and growth in the VLBW infant. Pediatr Res. 19: 1085, 1985.

97. Roy CC, Ste-Marie M, Chartrand L, Weber A, Bard H, Doray B. Correlation of the malabsorption of the premature infant with a medium-chain triglyceride formula. J Pediatr. 86: 446-50, 1975.

98. Gompertz S, Sammons H. The origin of fecal lipids: the composition of fecal fats in human subjects. Clin Chim Acta 8: 591-603, 1963.

99. Moya M, Domenech E. Role of calcium-phosphate ratio in milk formula on calcium balance in low birth weight infants during the first three days of life. Pediatr Res. 16: 675-81, 1982.

100. Widdowson Elsie M, Mc Cance R, Harrison CJ, Sutton A. Effects of giving phosphate supplements to breast-fed babies on absorption and excretion of calcium, strontium, magnesium and phosphorus. Lancet 2: 1250-1, 1963.

101. Barltrop D, Mole R, Sutton A. Absorption and endogenous fecal excretion of calcium by low birth weight infants on feed with varying contents of calcium and phosphate. Arch Dis Child. 52: 41-9, 1977.

102. Giles MM, Fenton MH, Shaw B, Elton RA, Clarke M, Lang M, Hume R. Sequential calcium and phosphorus balance studies in preterm infants. J Pediatr. 110 (4): 591-8, 1987.

103. Horsman A, Ryan SW, Congdon PJ, Truscott JG, Simpson M. Bone mineral accretion rate and calcium intake in preterm infants. Arch Dis Child. 64 (7): 910-8, 1989.

104. Younaszai M. Development of intestinal calcium transport. Textbook of Gastroenterology and Nutrition in Infancy. (E Lebenthal, ed.), Raven Press, New York, pp.623-9, 1981.

105. Pansu D, Bellaton C, Bronner F. Developmental changes in the mechanisms of duodenal calcium transport in the rat. Am J Physiol. 244: G20-6, 1983.

106. Senterre J, Salle B. Calcium and phosphorus economy of the preterm infant and its interaction with vitamin D and its metabolites. Acta Pediatr Scand Suppl. 296: 85-92, 1982.

107. Sann L, David L, Loras B, Lahet C, Frederich A, Bethenod M. Neonatal hypercalcemia in preterm infants fed with human milk. Helv Pediatr Acta 40 (2-3): 117-26, 1985.

108. Chan GM, Mileur L, Hansen JW. Calcium and phosphorus requirements in bone mineralization of preterm infants. J Pediatr. 113 (1 Pt 2): 225-9, 1988.

109. Bronner F, Pansu D, Stein WD. An analysis of intestinal calcium transport across the rat intestine. Am J Physiol. 250 (5 Pt 1): G561-9, 1986.

110. Schedl HP, Christensen K, Ronnenberg W, Thomas S. Intestinal calcium

transport in systemic disease. Studies with brush-border membrane vesicles. Mineral Electrolyte Metab. 16 (2-3): 147-53, 1990.

111. Sugimoto T, Ritter C, Slatopolsky E, Morrissey J. Effect of 1,25-dihydroxyvitamin D_3 on phospholipid metabolism in cultured bovine parathyroid cells. Endocrinology 122 (6): 2387-92, 1988.

112. Favus MJ. Factors that influence absorption and secretion of calcium in the small intestine and colon. Am J Physiol. 248: 9147-57, 1985.

113. Theofan G, King MW, Hall AK, Norman AW. Expression of calbindin-D28K mRNA as a function of altered serum calcium and phosphorus levels in vitamin D-replete chick intestine. Mol Cell Endocrinol. 54 (2-3): 135-40, 1987.

114. Hall AK, Norman AW. Regulation of calbindin-D28K gene expression in the chick intestine: effects of serum calcium status and 1,25-dihydroxyvitamin D_3. J Bone Miner Res. 5 (4): 331-6, 1990.

115. Meyer MH, Meyer RA Jr, Iorio RJ. A role for the intestine in the bone disease of juvenile X-linked hypophosphatemic mice: malabsorption of calcium and reduced skeletal mineralization. Endocrinology 115 (4): 1464-70, 1984.

116. Jirapinyo P, Srimaruta N, Lebenthal E, Lee PC. Vitamin D deficiency. pancreatic and small intestinal enzyme development in rats. Biol Neonate 54 (1): 20-8, 1988.

117. Salle B, David L, Glorieux F, Delvin E, Senterre J, Renaud H. Early oral administration of vitamin D and its metabolites in premature neonates on mineral homeostasis. Pediatr Res. 16: 75-8, 1982.

118. Salle BL, Senterre J, Glorieux FH, Delvin EE, Putet G. Vitamin D metabolism in preterm infants. Biol Neonate 52 Suppl. 1: 119-30, 1987.

119. Salle BL, Senterre J, Glorieux FH, Delvin EE, Putet G. Vitamin D metabolism in preterm infants. Biol Neonate 52 Suppl. 1: 119-30, 1987.

120. Boden SD, Kaplan FS. Calcium Homeostasis. Orthop Clin North Am. 21 (1)F: 31-42, 1990.

121. Williams MC, C.Morrow G, et al. Calcium and fat absorption in the neonatal period. Am J Clin Nutr. 23: 1322-30, 1970.

122. Atteh JO, Leeson S. Effects of dietary saturated or unsaturated fatty acids and calcium levels on performance and mineral metabolism of broiler chicks. Poult Sci. 63 (11)f: 2252-60, 1984.

123. Warshaw B, Anand S, Kerian A, Liberman E. The effect of chronic furosemide administration on urinary calcium excretion and calcium balance in growing rats. Pediatr Res. 14: 1118-21, 1980.

124. Brown D, Steranka B. Renal cation excretion in the hypocalcemic premature human neonate. Pediatr Res. 15: 1100-4, 1981.

125. Brannan P, Vergne-Marini P, Pak C, Hall A, Fordtran J. Magnesium absorption in the human small intestine. J Clin Invest. 57: 1412-8, 1976.

126. Venkataraman P, Han B, Tsang R, Danghterty C. Secondary hyperparathyroidism and bone disease in infants receiving long-term furosemide therapy. Am J Dis Child. 137: 1157-61, 1983.

127. Hufnagel K, Khan S, Penn D, Cucciarelli A, Williams P. Renal calcifications: a complication of long-term furosemide therapy in preterm infants. Pediatrics 70: 360-3, 1982.

128. Ghishan F, Meneely R. Intestinal maturation: the effect of glucocorticoids on in vivo net magnesium and calcium transport in the rat. Life Sci. 31: 133-8, 1982.

129. Ammann P, Rizzoli R, Fleisch H. Influence of the disaccharide lactitol on intestinal absorption and body retention of calcium in rats. J Nutr. 118 (6): 793-5, 1988.

130. Karlen J, Aperia A, Zetterstrom R. Renal excretion of calcium and phosphate in preterm and term infants. J Pediatr. 106 (5): 814-9, 1985.

131. Venkataraman PS, Lick KE. Effect of mineral supplementation of human milk on bone mineral content and trace element metabolism. J Pediatr. 113 (1 Pt 2): 220-4, 1988.

132. Greer FR, McCormick A. Improved bone mineralization and growth in premature infants fed fortified own mother's milk. J Pdiatr. 112 (6): 961-9, 1988.

133. Lyon AJ, McIntosh N. Calcium and phosphorus balance in extremely low birthweight infants in the first six weeks of life. Arch Dis Child. 59 (12): 1145-50, 1984.

134. Manz F, Diekmann L, Stock GJ. Effect of calcium supplementation on calcium and phosphorus balance and renal net acid excretion in preterm infants fed a standard formula. Acta Paediatr Scand. 78 (4): 525-31, 1989.

135. Gafter U, Edelstein S, Hirsh J, Levi J. Metabolic acidosis enhances

1,25 $(OH)_2D_3$-induced intestinal absorption of calcium and phosphorus in rats. Mineral Electrolyte Metab. 12 (4): 213-7, 1986.

136. Armbrecht HJ. Age-related changes in calcium and phosphorus uptake by rat small intestine. Biochim Biophys Acta 882 (3): 281-6, 1986.

137. Barrionuevo M, Campos MS, Lopez-Aliaga I, Coves F, Lisbona F. Nutritive utilization of phosphorus in the rat: influence of intestinal resection and dietary medium chain triglycerides and vitamin D_3. Int J Vitam Nutr Res. 59 (3): 255-61, 1989.

138. Spencer H, Kramer L. Osteoporosis: calcium, fluoride, and aluminum interactions. J Am Coll Nutr. 4 (1): 121-8, 1985.

139. Greger JL, Kaup SM, Behling AR. Calcium, magnesium and phosphorus utilization by rats fed sodium and potassium salts of various inorganic anions. J Nutr. 121: 1382-8, 1991.

140. Hardwick LL, Hones MR, Brautbar N, Lee DB. Magnesium absorption: mechanisms and the influence of vitamin D, calcium and phosphate. J Nutr. 121 (1): 13-23, 1991.

141. Brink EJ, Dekker PR, Van Beresteiju ECH, Beyuan AC. Inhibitory effect of dietary soybean protein vs casein on magnesium absorption rats. J Nutr. 121: 1374-81, 1991.

142. Hardwick LL, Hones MR, Buddington RK, Clemens RA, Lee DB. Comparison of calcium and magnesium absorption: in vivo and in vitro studies. Am J Physiol. 259 (5 Pt 1): G720-6, 1990.

143. Schneider KM, Parkinson GB, Houston JC, Leaver DD. 1,25 dihydroxyvitamin D_3 increases plasma magnesium and calcium in sheep fed liquid diets low in calcium and magnesium. Aust Vet J. 62 (3): 82-5, 1985.

144. Selby PL, Peacock M, Bambach CP. Hypomagnesaemia after small bowel resection: treatment with 1-alpha-hydroxylated vitamin D metabolites. Br J Surg. 71 (5): 334-7, 1984.

145. Giles MM, Laing IA, Elton RA, Robins JB, Sanderson M, Hume R. Magnesium metabolism in preterm infants: effects of calcium, magnesium, and phosphorus, and of postnatal and gestational age. J Pediatr. 117 (1): 147-54, 1990.

146. Rahnema SH, Fontenot JP. Effects of intravenous infusion of high levels of potassium and sodium on mineral metabolism in sheep. J Anim Sci. 68 (9): 2833-8, 1990.

147. Scharrer E, Lutz T. Effects of short chain fatty acids and K on

absorption of Mg and other cations by the colon and caecum. Z Ernahrungswiss. 29 (3): 162-8, 1990.

148. Aliaga IL, Miller DL, Wilson HD, Schedl HP. Effects of resection on absorption and secretion of divalent cations by small intestine of rat. Am J Clin Nutr. 52 (5): 867-71, 1990.

149. Caddell JL. Protection by magnesium of renal calcinosis in furosemide-treated weanling rats with moderate magnesium deficiency. Biol Neonate 48 (1): 49-58, 1985.

Chapter II

CALCIUM SUPPLEMENTATION IN PREMATURE INFANTS FED BREAST MILK: The Effect of Timing of Calcium Supplementation on Calcium, Phosphorus, Magnesium and Fat Balance and on the Absorption Coefficients of Individual Fatty Acids

A: Introduction

Lipid and calcium soap formation has been extensively studied in newborn infants by giving various forms of calcium supplement with either human milk or formulas. However, giving the supplement other than together with feedings has not been studied. Therefore, in this study, sixteen medically stable premature newborns were pair-matched according to their gestational age and birthweight, divided into two groups at random and given calcium lactate either with or between expressed human milk feedings in the first four weeks of life. The measurement and observation focused on growth parameters, lipid, calcium (Ca), phosphorus (P) and magnesium (Mg) balance and the absorption coefficients of certain fatty acids. This chapter will present the experimental design, methodology, results of the study by tables and figures and discussion of these outcomes.

B: Study Subjects and Methodology

Infant nursing, milk and supplement feeding, sample collecting and anthropometric measuring were carried out by the research nurse in NICU. Calcium, phosphorus and magnesium in milk, stool, and urine and the total fat in milk and stool were measured in the Neonatal Research Laboratory of University of Alberta.

1. Study Design

Study Subject Description

Three-day metabolic balance studies were carried out during the first, second and fourth postnatal week in 16 infants born at 30.8±1.4 weeks gestation and weighing 1329±168 grams (Mean±SD) at birth at the Neonatal Intensive Care Unit (NICU) of the University of Alberta Hospital. All infants had normal renal function and acid-base balance, no congenital abnormalities, no steroid, diuretic or sodium medications and tolerated full feeds. Infants were nursed in incubators in a thermoneutral environment (1) (infants' core temperature ranged from 36.7 °C to 37.3 °C).

Feeding Regimen

Infants were pair-matched according to gestational age and birthweight and randomly divided into two groups. After expressed own mother's milk was collected (see Collection of Preterm Human Milk), infants in both groups were fed with calibrated syringes every two hours.

Calcium Supplementation

Once the infants tolerated 50% oral feeds, calcium lactate (1.3 mmole calcium /Kg /day) was administered every two hours with milk to 8 infants and exactly in between feedings to the other 8 infants. Other oral supplements, including vitamin C (100mg for first week), vitamin E (50IU for 60 days or until discontinuation of oxygen, whichever was longer), and vitamin D (1000IU), were given daily to each infant.

2. Collection of Preterm Human Milk

Expressed own mother's milk was delivered fresh or frozen to the NICU within 24 hours after collection in sterile glass containers. The milk was mixed and then divided into aliquots for feeding from the total 24-hour expressed volume. A 10 ml aliquot for each study day was frozen at -20 °C for analysis of calcium, phosphorus, magnesium, total fat and fatty acids. All expressed breast milk was fed within 48 hours of fresh collection or within 24 hours after thawing. Volume of milk intake was calculated from the fluid fed with calibrated syringes, minus gastric aspirates and vomitus which, if it occurred, was collected on ashless paper placed beneath the infant's mouth.

3. Collections of Urine and Stool Samples

Samples were collected, pooled and volumes measured in each study period. Urine was collected with an adaptive urine collector. All urine samples from each balance period were pooled and the total volume measured. An aliquot was removed and stored at -20 °C for later analysis.

Leakage was calculated from the weight of a pre-weighed diaper. Stools were collected on a large square sheet of polyethylene placed under the infant, frozen immediately at -20°C. They were later pooled and weighed for the 72-hour period, and then freeze dried and stored at -20°C for analysis.

4. Anthropometric Measurement

Anthropometric measurements included weekly head circumference using a fiberglass measuring tape, weekly body length using a plexiglass measuring board calibrated in milimeter increments, weekly skinfold at the triceps, subscapular and periumbilical regions with a Harpenden caliper stabilized for 30 seconds and daily body weight with an electronic scale to an accuracy of ± 1 gram.

5. Calcium, Phosphorus, Magnesium, Total Fat and Fatty Acids Measurement

Calcium in milk and supplement, stool and urine was measured with atomic absorption flame spectrophotometry (2). Lyophilised stool sample was wet ashed, and urine was diluted before calcium analysis. Phosphorus in milk, stool and urine was measured with a modified method of Fiske and Subbarow (3). Magnesium in all milk, stool and urine samples was also measured by atomic absorption spectrophotometry (2). Total fat in milk was measured by a gravimetric method (4), and fatty acids in milk were extracted with a modified Folch extraction (5) (see Appendix I) and analyzed by capillary gas-liquid chromatography (6). Total fat in stool was measured by a modified Jeejeebhoy procedure with the addition of HCl

to release bound fatty acids (7). Fatty acids in stool were extracted by a modified method of Watkins and Bliss in triplicate (8) (see Appendix II) and then analyzed by capillary gas-liquid chromatography (6). Blood work included initial acid-base status, calcium and phosphorus levels, creatinine and blood urea nitrogen. Both calcium and phosphorus levels were measured at the end of the study as well.

6. Statistics Applied

Statistics include multiple analysis of variance and correlation analysis. In addition, Student-Newman-Keuls test was used to further compare the difference between three study weeks. Student t test was used to measure the difference between the two treatments within each study week.

C: Results

An overall significant variation among individual premature infants was observed. Results are presented in tables and figures. Data are expressed as means ± SD.

1. Composition of Preterm Human Milk

Table II-1 shows the composition of milk consumed by the study subjects. Infants from different groups ingested similar volumes of milk per kilogram body weight per day. The pooled intake volume was 162±17 ml/Kg/d. Solid weight and total lipid content of the milk were the same for the different groups and different study weeks (pooled value: 12.6±0.7

Table II-1: Composition of Expressed Preterm Human Milk Consumed by Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=7	n=6
Volume (ml/Kg/d)			
Ca between	153±16	159±15	164±15
Ca with	164±18	163±19	172±19
Solid Weight (g/dl)			
Ca between	12.3±0.4	13.0±0.8	12.4±0.7
Ca with	12.7±0.6	12.9±0.6	12.5±0.9
Total Lipid (g/dl)			
Ca between	3.05±0.44	3.72±0.55	3.39±0.87
Ca with	3.41±0.62	3.33±0.35	3.16±0.91
Calcium (mmole/dl)*			
Ca between	0.68±0.19	0.60±0.16	0.55±0.14
Ca with	0.63±0.14	0.56±0.13	0.60±0.09
Phosphorus (mmole/dl)**			
Ca between	0.54±0.11	0.50±0.12	0.40±0.06
Ca with	0.51±0.11	0.48±0.06	0.47±0.07
Magnesium (mmole/dl)			
Ca between	0.14±0.05	0.14±0.05	0.12±0.04
Ca with	0.13±0.02	0.12±0.02	0.12±0.04

1. Data are presented as Mean ± SD;

2. * significant difference among postnatal weeks ($P<0.05$);

3. ** significant difference among postnatal weeks ($P<0.01$).

and 3.3±0.6 g/dl, respectively). There was no difference in Ca, P, and Mg content between the two groups. However, a progressive decline of Ca and P concentrations with postnatal age was observed. Preterm human milk had a significantly higher concentration of Ca in the first week after delivery which decreased gradually during the following study weeks (0.65±0.17, 0.58±0.14, 0.57±0.11 mmole/dl, respectively). The high P content in the milk lasted for the first two weeks of life (0.53±0.11, 0.49±0.09, 0.43±0.07 mmole/dl). However, the longitudinal decline in Mg concentration was not significant (0.13±0.04, 0.13±0.04, 0.12±0.04 mmole/dl).

2. Anthropometric Data

Table II-2 shows anthropometric data of the study subjects. The postnatal age and body weight at which the balance studies were conducted were not different between the two treatment groups. The growth parameters such as body weight, head circumference and length increased progressively with postnatal age and no difference was observed between the two treatments. Infants receiving calcium supplement between feedings showed less weight gain when compared to infants receiving calcium with feedings, especially in the second postnatal week (one-tailed t test: $P<0.05$), although it was not statistically significant. The pooled weight gain was 11.8±4.0 g/Kg/d. The mean weight gains in 'Ca between feedings' and 'Ca with feedings' were 11.4±4.6 and 12.2±4.1 g/Kg/d, respectively. The mean weight gains during three study periods were 9.7±3.9, 11.7±4.1, and 13.9±5.3 g/Kg/d, respectively. However, none of these was

Table II-2: Anthropometric Data for Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=7	n=6
Age (gestational week) *			
Ca between	32.1±1.3	33.3±1.3	34.7±0.8
Ca with	31.9±1.7	33.3±1.4	34.0±1.4
Weight (g) *			
Ca between	1276±195	1450±211	1598±358
Ca with	1300±201	1461±223	1540±239
Head Circumference (cm) *			
Ca between	28.2±1.9	29.6±1.8	30.2±2.0
Ca with	28.0±1.8	29.0±1.8	29.9±1.9
Body Length (cm) *			
Ca between	40.0±2.4	41.8±1.0	41.9±3.1
Ca with	39.8±1.9	41.4±1.8	41.8±1.7
Weight Gain (g/Kg/d)			
Ca between	9.7±3.3	9.8±3.0	15.3±5.8
Ca with	10.7±3.1	13.6±4.4	12.5±4.8

1. Data are presented as Mean ± SD;

2. * significant difference among postnatal weeks ($P<0.01$).

statistically significant.

3. Balance Studies

In general, balance studies show increased fecal excretions of lipid, Ca, P, and Mg in the infants receiving Ca between feedings compared to the infants receiving Ca with feedings. The absorption and retention rate of these elements was correspondingly affected. The following summaries and tables present detailed outcomes of each balance study.

Lipid Balance Study

As milk intake and lipid content of the milk were relatively consistent, lipid intakes between groups during the three balance studies were quite similar. The pooled value was 5.40±1.11 g/Kg/d. The lipid loss in stools was higher in infants given Ca between feedings compared to infants given Ca with feedings, but the difference was not statistically significant (1.30±0.85 vs 0.95±0.80 g/Kg/d). However, the difference between lipid intake and lipid loss in stool was statistically significant in the first postnatal week (two-tailed t test: $P<0.05$). This may, in part, explain the difference in total lipid absorption rates between the two groups (75.4±16.8% vs 82.3±15.3%), which was not statistically significant (Table II-3).

Calcium Balance Study

There was no difference in Ca intake between groups. Infants from both groups ingested significantly more Ca in the first week of life,

Table II-3: Lipid Balance in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=7	n=6
Fat intake (g/Kg/d)			
Ca between	4.72±1.10	5.91±0.96	5.52±1.33
Ca with	5.56±1.08	5.45±1.06	5.34±1.15
Fat loss in stool (g/Kg/d)			
Ca between	1.19±0.86	1.42±1.00	1.29±0.77
Ca with	0.92±0.75	0.87±0.84	1.08±0.93
Fat intake - Fat loss in stool (g/Kg/d)			
Ca between	3.53±0.50 *	4.49±1.16	4.23±1.89
Ca with	4.64±1.06 *	4.57±0.78	4.26±1.79
Fat absorption (%)			
Ca between	76.7±11.6	76.0±16.2	73.0±24.8
Ca with	83.5±13.1	85.1±12.6	77.5±21.6

1. Data are presented as Mean ± SD;
2. * significant difference between two treatments during a study week ($P<0.05$).

Table II-4: Calcium Balance in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1 n=8	Week 2 n=7	Week 4 n=6
Ca intake from milk (mmole/Kg/d)			
Ca between	1.03±0.26	0.96±0.27	0.89±0.23
Ca with	1.04±0.30	0.92±0.27	1.03±0.14
Ca Lactate intake (mmole/Kg/d) *			
Ca between	1.36±0.23	1.18±0.08	1.08±0.18
Ca with	1.35±0.23	1.21±0.10	1.14±0.16
Total Ca intake (mmole/Kg/d) *			
Ca between	2.29±0.39	2.13±0.26	1.97±0.25
Ca with	2.31±0.31	2.13±0.32	2.17±0.23
Ca loss in stool (mmole/Kg/d)			
Ca between	0.87±0.52	0.71±0.46	0.65±0.25
Ca with	0.67±0.36	0.50±0.29	0.58±0.39
Ca absorption (%)			
Ca between	63.7±21.4	66.4±22.9	67.5±10.4
Ca with	71.5±12.4	77.4±10.3	73.9±17.0
Ca loss in urine (mmole/Kg/d) *			
Ca between	0.48±0.31	0.56±0.19	0.62±0.20
Ca with	0.38±0.24	0.54±0.23	0.61±0.25
Total Ca loss (mmole/Kg/d)			
Ca between	1.28±0.53	1.27±0.41	1.27±0.21
Ca with	1.05±0.32	1.06±0.16	1.19±0.35
Ca retention (%)			
Ca between	46.8±19.8	40.3±20.3	35.8±5.3
Ca with	55.1±10.7	51.1±5.2	45.3±14.7

1. Data are presented as Mean ± SD;
2. * significant difference among postnatal weeks ($P<0.05$).

because of the higher intake of Ca from the supplement during the first balance study and the higher Ca content in human milk. Pooled values of Ca total intake during the three studies were 2.35±0.35, 2.13±0.28, 2.07±0.25 mmole/Kg/d, respectively. Infants receiving Ca supplement between feedings excreted more Ca in the stool compared to infants receiving Ca supplement with feedings (0.76±0.43 vs 0.59±0.34 mmole/Kg/d). This accounted for the increased total Ca loss in infants receiving Ca between feedings (1.27±0.40 vs 1.09±0.28 mmole/Kg/d). The increased Ca loss contributed to the impaired absorption (65.7±18.7% vs 74.1±12.8%) and retention rates (41.5±17.0% vs 51.0±10.9%) in infants receiving Ca supplement between feedings, although none of these values was statistically significant. No significant difference between groups was observed in Ca loss in urine. Infants from both groups excreted more Ca in urine after the first week of life (0.43±0.28, 0.55±0.20 and 0.61±0.22 mmole/Kg/d from the first week to the fourth week, respectively). This change paralleled the change in Ca retention rate (50.9±15.9%, 45.7±15.3%, 40.5±11.6% from the first week to the fourth week, respectively), although the latter was not statistically significant (Table II-4).

Phosphorus Balance Study

In the P balance study, infants from both groups consumed significantly more P in the first week of life than the fourth week (0.83±0.15 vs 0.73±0.13 mmole/Kg/d). More P was excreted from stools in infants given Ca between feedings (0.11±0.09 vs 0.06±0.04 mmole/Kg/d) and total P loss was increased as well (0.11±0.09 vs 0.06±0.04 mmole/Kg/d),

Table II-5: Phosphorus Balance in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=7	n=6
P intake (mmole/Kg/d) *			
Ca between	0.82±0.14	0.80±0.18	0.66±0.09**
Ca with	0.83±0.17	0.78±0.16	0.81±0.13**
P loss in stool (mmole/Kg/d)			
Ca between	0.13±0.13	0.09±0.05	0.10±0.05
Ca with	0.06±0.03	0.06±0.04	0.06±0.04
P absorption (%) **			
Ca between	85.6±13.1	88.2±8.0	84.4±7.1**
Ca with	93.2±4.0	92.9±4.3	92.9±4.7**
P loss in urine (mmole/Kg/d)			
Ca between	0.003±0.003	0.002±0.002	0.002±0.002
Ca with	0.005±0.006	0.004±0.002	0.006±0.008
Total P loss (mmole/Kg/d)			
Ca between	0.13±0.13	0.09±0.05	0.11±0.05
Ca with	0.06±0.04	0.06±0.04	0.06±0.04
P retention (%) **			
Ca between	85.3±12.9	87.9±8.0	84.0±7.1**
Ca with	92.5±4.3	92.4±4.3	92.5±4.9**

1. Data are presented as Mean ± SD;

2. * significant difference among postnatal weeks ($P<0.05$);

3. ** significant difference between two treatments ($P<0.05$).

although they were not statistically significant. This increased P loss in stools resulted in a significant decrease in both P absorption (86.1±9.7% vs 93.0±4.1%) and retention rates (85.8±9.6% vs 92.5±4.3%). Unexpectedly, t test shows that infants receiving Ca between feedings consumed more P in the fourth week (two-tailed t test: $P<0.05$), and both P absorption and retention in this week were significantly increased (two-tailed t test: $P<0.05$). P level in urine was minimal and was similar in both groups during the three weeks. Pooled mean value of P loss in urine was 0.003±0.003 mmole/Kg/d (Table II-5).

Magnesium Balance Study

The average Mg intake was 0.20±0.06 mmole/Kg/d. No difference was found between groups or study weeks. Fecal Mg excretion increased in infants receiving Ca between feedings (0.10±0.07 vs 0.06±0.03 mmole/Kg/d), although the increase was not significant. Total Mg loss was, however, significantly increased in these infants (0.18±0.06 vs 0.13±0.02 mmole/Kg/d). Mg absorption (50.3±32.4% vs 68.7±12.4%) and retention rates (6.4±33.4% vs 33.2±14.2%) were altered as well, although only the latter was statistically significant. The difference of Mg retention in the first week was statistically significant in two-tailed t test ($P<0.05$). A progressive decline in Mg in urine (0.08±0.03, 0.08±0.02 and 0.06±0.04 mmole/Kg/d, respectively) and total Mg loss (0.17±0.07, 0.15±0.04 and 0.13±0.03 mmole/Kg/d, respectively) during the study were also recorded. However, the increased Mg retention rate with postnatal age was not statistically significant (16.0±32.4%, 19.9±26.7% and 24.7±27.6%,

Table II-6: Magnesium Balance in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=7	n=6
Mg intake (mmole/Kg/d)			
Ca between	0.21±0.07	0.23±0.10	0.19±0.06
Ca with	0.21±0.03	0.19±0.04	0.20±0.07
Mg loss in stool (mmole/Kg/d)			
Ca between	0.12±0.09	0.09±0.05	0.08±0.04
Ca with	0.07±0.03	0.06±0.04	0.06±0.03
Mg absorption (%)			
Ca between	42.2±41.2	54.9±30.6	55.7±23.2
Ca with	68.3±13.4	69.6±15.3	68.2±9.1
Mg loss in urine (mmole/Kg/d) *			
Ca between	0.09±0.03	0.08±0.02	0.08±0.04
Ca with	0.07±0.02	0.07±0.02	0.05±0.03
Total Mg loss (mmole/Kg/d) *,**			
Ca between	0.21±0.08**	0.18±0.04**	0.15±0.03**
Ca with	0.13±0.02**	0.13±0.02**	0.12±0.03**
Mg retention (%) **			
Ca between	-3.0±36.6**	11.7±36.6	12.5±27.7
Ca with	35.0±9.4**	28.0±7.2	36.8±23.7

1. Data are presented as Mean ± SD;

2. * significant difference among postnatal weeks ($P<0.01$);

3. ** significant difference between two treatments ($P<0.05$).

respectively) (Table II-6).

4. Correlation of Calcium Parameters

Figure II-1 shows the correlation analysis of Ca intake and Ca loss in stool ($P<0.01$). Ca intake positively correlated with Ca loss in stools, this correlation being more obvious in infants receiving Ca with feeding ($r=0.70$ vs $r=0.33$).

5. Correlations of Fecal Lipid, Calcium, Phosphorus and Magnesium

Figures II-2 to II-6 illustrate the result of fecal total lipid, Ca, P, and Mg correlation analysis ($P<0.01$). Fecal lipid was significantly correlated with Ca and P content in stools. The correlations were greater in infants given the Ca supplement with milk (lipid and Ca: $r=0.91$ vs $r=0.63$; lipid and P: $r=0.92$ vs $r=0.23$). Fecal P and Mg were strongly correlated in both groups ($r=0.77$ and $r=0.78$). In the 'Ca with feeding' group the correlation between Ca and P was very obvious ($r=0.83$), while in the other group the correlation between Ca and Mg was more significant ($r=0.82$).

6. Absorption Coefficients of Individual Fatty Acids

Table II-7 shows the absorption coefficient of fatty acids C12:0, C14:0, C16:0, C18:0, C18:1(9) and C18:2(6). The absorption coefficients of C14:0, C16:0, C18:0, and C18:1(9) appear to decrease in 'Ca between feedings' group, especially in the first two weeks, although they were not statistically significant (C14:0-81.4±20.7% vs 91.3±8.1%, C16:0-72.1

Figure II-1: Correlation of Ca Intake and Ca Loss in Stool in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week



Figure II-2: Correlation of Fecal Lipid and Ca Losses in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week



Figure II-3: Correlation of Fecal Lipid and P Losses in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week



Figure II-4: Correlation of Fecal P and Mg losses in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second and Fourth Postnatal Week



Figure II-5: Correlation of Fecal Ca and P Losses in Breast Fed Premature Infants Receiving Ca Lactate With Feedings in the First, Second and Fourth Postnatal Week



Figure II-6: Correlation of Fecal Ca and Mg Losses in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings in the First, Second and Fourth Postnatal Week



Table II-7: Absorption Coefficients for Selected Fatty Acids in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=6	n=6
C12:0 (%)			
Ca between	98.3±3.8	94.8±10.4	100.0±0.0
Ca with	99.5±0.9	99.3±1.3	98.9±2.7
C14:0 (%)			
Ca between	84.9±14.7	69.7±30.3	88.2±12.9
Ca with	91.2±8.1	93.2±6.0	89.7±10.8
C16:0 (%)			
Ca between	74.6±25.8	56.1±40.6	84.8±10.2
Ca with	85.9±11.8	87.5±10.4	84.8±17.8
C18:0 (%)			
Ca between	52.8±30.2	33.9±40.9 *	57.6±13.3
Ca with	71.2±16.6	75.6±17.0 *	71.5±23.9
C18:1(9) (%)			
Ca between	73.7±24.5	60.7±35.5	75.9±20.9
Ca with	76.5±27.1	87.5±13.5	77.3±26.8
C18:2(6) (%)			
Ca between	85.9±10.9	70.1±36.4	68.3±36.5
Ca with	69.6±38.7	78.0±38.8	72.1±39.3

1. Data are presented as Mean ± SD;

2. * significant difference between two treatments ($P<0.05$).

±29.0% vs 86.0±12.8%; C18:0-48.6±30.4% vs 72.6±18.2%; C18:1(9)-70.5±26.7% vs 80.1±23.1%). Only the absorption coefficient of fatty acid C18:0 in the infants receiving Ca between feedings in the second study week was significantly different from that in the infants receiving Ca with feedings (two-tailed t test: $P<0.05$).

Table II-8 illustrates the quantitative measurement of these six fecal fatty acids. Infants receiving Ca between feedings excreted more C14:0, C16:0, C18:0, C18:1(9) and C18:2(6) in the first two postnatal weeks compared to infants receiving Ca with feedings (C14:0-0.07±0.06 vs 0.05±0.06g/Kg/d; C16:0-0.30±0.23 vs 0.25±0.28g/Kg/d; C18:0-0.17±0.08 vs 0.13±0.12g/Kg/d; C18:1(9)-0.40±0.35 vs 0.32±0.34g/Kg/d; C18:2(6)-0.07±0.07 vs 0.04±0.05g/Kg/d). However, these differences were also not statistically significant.

7. Fecal Lipid Fractions

Table II-9 presents the percentage of four fecal lipid fractions: neutral lipid, Ca salts, free fatty acids, and compound lipid (the pooled values were approximately 54.4%, 2.5%, 21.5%, and 21.7%, respectively). Neutral lipid excretion increased gradually in the first four weeks of life (48.6±21.1%, 54.3±20.7%, and 62.1±17.7%, respectively), whereas free fatty acids decreased during the same period (25.8±13.5%, 19.7±10.9%, and 17.5±7.5%, respectively). In infants given Ca between feedings, the percentage of free fatty acids in stool was decreased (18.2±9.9% vs 24.8±12.3%) and the percentage of neutral lipid increased (57.1±23.7% vs 51.6±16.3%), although they were not statistically significant.

Table II-8: Quantitative Measurement of Selected Fecal Fatty Acids in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=6	n=6
C12:0 (mg/Kg/d)			
Ca between	5±11	13±25	0±0
Ca with	1±2	6±14	9±21
C14:0 (mg/Kg/d)			
Ca between	63±50	97±72	41±44
Ca with	48±50	49±54	64±81
C16:0 (mg/Kg/d)			
Ca between	302±211	419±290	180±128
Ca with	236±239	219±213	286±402
C18:0 (mg/Kg/d)			
Ca between	165±77	204±97	145±35
Ca with	125±79	120±93	156±179
C18:1(9) (mg/Kg/d)			
Ca between	382±352	493±402	312±348
Ca with	293±327	308±391	356±372
C18:2(6) (mg/Kg/d)			
Ca between	61±66	88±82	52±68
Ca with	37±46	40±57	52±67

Data are presented as Mean ± SD.

Table II-9: Percentages of Fecal Lipid Fractions in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=6	n=6
Neutral Lipid (%) *			
Ca between	51.2±26.5	59.5±24.4	62.7±21.4
Ca with	46.2±15.4	49.2±16.8	61.4±15.2
Ca Salts (%)			
Ca between	2.9±2.5	1.6±0.6	3.2±3.2
Ca with	2.6±1.9	2.3±2.2	2.0±1.4
Free Fatty Acids (%) *			
Ca between	21.8±11.9	17.2±10.2	14.2±5.7
Ca with	29.8±14.6	22.2±12.0	20.8±8.1
Compound Lipid (%)			
Ca between	24.2±21.2	21.9±16.5	19.9±22.8
Ca with	21.5±11.8	26.1±15.9	15.7±9.8

1. Data are presented as Mean ± SD;

2. * significant difference among postnatal weeks ($P<0.05$).

Data of fecal lipid fractions from Table II-10 shows the quantitative change in these fractions. Table II-11a to Table II-11f show the fractional distribution of selected fatty acids in stool. In general, more neutral lipid that consisted of fatty acid C14:0, C16:0, and C18:0 had been hydrolyzed (65% of fecal fatty acid C14:0 was found in neutral lipid fraction, 47% of C16:0, and 33% of C18:0 vs 83% of C12:0, 86% of C18:1, and 91% of C18:2). Infants receiving Ca between feedings seem to have excreted more C16:0, C18:0, C18:1(9), and C18:2(6) in the form of neutral lipid (C16:0-156±187 vs 98±121 mg/Kg/d; C18:0-62±53 vs 38±38 mg/Kg/d; C18:1(9)-351±356 vs 246±271 mg/Kg/d; C18:2(6)-62±70 vs 34±44 mg/Kg/d). Longitudinal decline of free fatty acid formation also appears to exist in fatty acid C14:0 and C18:1(9) (C14:0-11.7±9.3, 9.6±6.4, 6.6±7.9 mg/Kg/d; C18:1(9)-33.1±42.1, 19.3±19.3, 15.2±11.1 mg/Kg/d). However, because of the great individual variations, none of these were statistically significant.

Table II-10: Quantitative Measurement of Fecal Lipid Fractions in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=6	n=6
Neutral Lipid (mg/Kg/d)			
Ca between	776±850	1179±1055	693±649
Ca with	484±517	493±540	746±813
Ca Salts (mg/Kg/d)			
Ca between	29±21	22±10	31±31
Ca with	27±34	27±30	26±36
Free Fatty Acids (mg/Kg/d)			
Ca between	238±154	228±132	118±33
Ca with	245±250	143±112	216±292
Compound Lipid (mg/Kg/d)			
Ca between	202±135	271±202	162±162
Ca with	166±147	290±482	174±265

Data are presented as Mean ± SD.

Table II-11a: Quantitative Measurement of Fecal Fatty Acid C12:0 in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=6	n=6
Neutral Lipid (mg/Kg/d)			
Ca between	4.39±9.75	11.89±24.05	0.00±0.00
Ca with	0.73±1.68	4.03±8.94	7.53±17.75
Ca Salts (mg/Kg/d)			
Ca between	0.03±0.05	0.08±0.12	0.00±0.00
Ca with	0.01±0.02	0.06±0.15	0.12±0.29
Free Fatty Acids (mg/Kg/d)			
Ca between	0.41±0.80	0.49±0.76	0.00±0.00
Ca with	0.34±0.69	0.58±1.19	0.54±1.22
Compound Lipid (mg/Kg/d)			
Ca between	0.11±0.24	0.63±1.24	0.00±0.00
Ca with	0.08±0.18	1.60±3.78	0.66±1.53

Data are presented as Mean ± SD.

Table II-11b: Quantitative Measurement of Fecal Fatty Acid C14:0 in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=6	n=6
Neutral Lipid (mg/Kg/d)			
Ca between	39.1±48.9	70.9±70.3	28.5±41.7
Ca with	26.6±35.6	27.0±30.2	45.8±57.4
Ca Salts (mg/Kg/d)			
Ca between	1.38±0.89	1.20±0.59	1.29±1.99
Ca with	1.86±2.95	1.58±1.87	1.52±2.28
Free Fatty Acids (mg/Kg/d)			
Ca between	12.0±7.7	12.1±6.0	4.3±1.9
Ca with	11.4±11.3	7.1±6.4	8.8±11.0
Compound Lipid (mg/Kg/d)			
Ca between	10.6±6.8	12.2±6.8	6.5±7.4
Ca with	7.7±7.1	13.5±21.6	7.9±12.2

Data are presented as Mean ± SD.

Table II-11c: Quantitative Measurement of Fecal Fatty Acid C16:0 in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=6	n=6
Neutral Lipid (mg/Kg/d)			
Ca between	141±183	237±248	95±106
Ca with	80±99	83±88	138±177
Ca Salts (mg/Kg/d)			
Ca between	11±9	8±5	9±13
Ca with	11±15	11±13	10±17
Free Fatty Acids (mg/Kg/d)			
Ca between	88±62	90±59	33±11
Ca with	89±102	52±41	88±142
Compound Lipid (mg/Kg/d)			
Ca between	61±52	84±53	43±47
Ca with	55±55	73±95	50±75

Data are presented as Mean ± SD.

Table II-11d: Quantitative Measurement of Fecal Fatty Acid C18:0 in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=6	n=6
Neutral Lipid (mg/Kg/d)			
Ca between	51.9±57.4	82.4±69.5	55.1±21.5
Ca with	28.7±24.7	33.0±23.1	54.9±59.4
Ca Salts (mg/Kg/d)			
Ca between	7.6±6.5	4.9±2.4	8.2±7.6
Ca with	5.7±5.9	5.8±5.8	6.4±8.5
Free Fatty Acids (mg/Kg/d)			
Ca between	59.4±34.9	60.2±36.3	37.9±16.6
Ca with	56.8±44.0	37.2±23.2	57.7±69.0
Compound Lipid (mg/Kg/d)			
Ca between	46.1±30.7	56.2±30.8	43.6±43.7
Ca with	33.7±21.9	56.8±81.0	36.5±45.8

Data are presented as Mean ± SD.

Table II-11e: Quantitative Measurement of Fecal Fatty Acid C18:1(9) in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=6	n=6
Neutral Lipid (mg/Kg/d)			
Ca between	332±352	449±406	278±354
Ca with	229±260	207±258	309±333
Ca Salts (mg/Kg/d)			
Ca between	3±2	4±2	4±5
Ca with	4±4	4±5	4±6
Free Fatty Acids (mg/Kg/d)			
Ca between	28±20	18±11	12±6
Ca with	39±58	20±26	18±15
Compound Lipid (mg/Kg/d)			
Ca between	20±18	22±16	18±21
Ca with	21±28	12±9	25±41

Data are presented as Mean ± SD.

Table II-11f: Quantitative Measurement of Fecal Fatty Acid C18:2(6) in Breast Fed Premature Infants Receiving Ca Lactate Between Feedings or With Feedings in the First, Second, and Fourth Postnatal Week

	Week 1	Week 2	Week 4
	n=8	n=6	n=6
Neutral Lipid (mg/Kg/d)			
Ca between	56.6±65.6	83.7±83.1	47.9±68.0
Ca with	30.3±35.7	22.7±28.2	50.6±64.8
Ca Salts (mg/Kg/d)			
Ca between	0.38±0.32	0.36±0.27	0.65±0.77
Ca with	0.49±0.73	0.54±1.10	0.32±0.30
Free Fatty Acids (mg/Kg/d)			
Ca between	2.53±2.25	1.46±0.96	1.47±0.92
Ca with	3.34±5.54	1.21±1.55	1.01±0.63
Compound Lipid (mg/Kg/d)			
Ca between	1.78±1.78	1.95±1.54	1.74±2.10
Ca with	3.04±6.17	4.22±8.81	1.55±2.07

Data are presented as Mean ± SD.

D: Discussion and Conclusions

1. Chronological Changes in Milk Composition

Calcium and phosphorus content of preterm milk decreased gradually in the first four weeks of life, which is consistent with previous reports (9,10,11,12). Mean calcium concentration of preterm milk during this period is also consistent with other studies (13,14). However, the decline in magnesium concentration was not as obvious as that in calcium and phosphorus. Mean milk volume consumed by infants from both groups in the first four postnatal weeks was 162 ± 17 ml/Kg/d which is within the range of 159-182 ml of other four similar studies (12,14,15,16). This volume may provide the infant with about 105 Kcal/Kg/d if it was well absorbed (16). The total lipid in preterm milk, 3.34 ± 0.63 g/dl, was a little less than the approximate amount of 4 g/dl in another study of preterm human milk (13).

2. Weight Gains in Anthropometric Study

Infants from both groups grew steadily as measured by body weight, body length and head circumference. However, weight gain of all infants in this study was less than other studies of non-supplemented preterm milk fed premature infants (14,16). Morever, infants receiving calcium between feedings showed even less weight gain compared to infants receiving calcium with feedings, especially during the second week of life, possibly because of the increased loss of ingested lipid from stool in relation to the decreased energy intake. Attendency to increase weight gain with

postnatal age in the first four weeks of life also seems to exist.

3. Effects of Calcium Supplementation on Intestinal Fat and Mineral Absorption

Mean lipid intake was about 5.4 ± 1.1 g/Kg/d which is in agreement with previous studies (12,15,16). Fecal lipid excretion in the infants given calcium between feedings was much higher when compared to the premature infants who were fed exclusively preterm human milk (15,16). The latter usually exhibits a fat absorption rate of more than 80% (15,17). However, the infants given calcium between feedings in the present study absorbed approximately 75% of milk lipid. Accordingly, all calcium, phosphorus, and magnesium absorption seem impaired when the calcium supplement was given between feedings with human milk (calcium 66% vs 74%, phosphorus 86% vs 93%, and magnesium 50% vs 72%).

In the original hypothesis, it is assumed that the less time that calcium and lipid interact in the intestinal lumen, the less chance they will have to form insoluble soap. However, the results of this experiment suggest that, as compared to giving calcium between feeding, giving calcium with the milk actually facilitates calcium and fat absorption, resulting in reduction of calcium-soap formation, and relatively less calcium and fat excretion in stool. As the interaction of minerals is also present in intestinal lumen, the change in fat absorption may alter phosphorus and magnesium absorption as well.

Possible Activation of Lipolytic Enzyme(s)

Some mechanisms may be involved in the enhancement of fat absorption when fat is ingested with calcium. Calcium may improve fat absorption by forming micelles, which incorporate or remove the lipolytic product of fat digestion, i.e. free fatty acids, in the stomach and intestine, thereby preventing a product inhibition effect. Another possibility is that calcium given with feedings activates the lipolytic enzymes, lipase(s), in the intestine. Calcium may have acted on pancreatic lipase or bile salt-stimulated lipase in the breast milk. The activated lipase digested milk fat more efficiently and, therefore, both fat and calcium absorption were enhanced. In vitro, calcium ion incubation decreases not only the lag phase before the accelerated hydrolysis of triglycerides reaches a high rate (18), but also decreases the rate of resynthesis of glyceride ester bonds (19). Calcium ions may have also acted on colipase synthesis and thus the binding strength and enzyme activity of lipase-colipase increases. In the presence of calcium ions, a smaller form of colipase, $colipase_{85}$, appears, whose specific activity is five times higher than that of regular $colipase_{101}$ (20). With intralipid as substrate, $colipase_{85}$ enables lipase to reach the triacylglycerol substrate more rapidly and shortens the lag phase for a given concentration (20). This type of interaction has further been confirmed recently by a study of triglyceride emulsions in vitro showing that the micelle-lipase-colipase complex exhibits decreased activity in the absence of calcium ions; whereas, in the presence of calcium, it becomes more active and binds more easily to interfaces (21).

Calcium Soap Formation

As expected, calcium soap formation occurred as shown from the positive correlation between fecal lipid and calcium content in the stool, especially in the infants receiving calcium with feedings (calcium with feedings: r=0.91 vs calcium between feedings: r=0.63). The long-chain saturated fatty acids were most involved in the soap formation and consequently excreted, as indicated by the increased loss of fatty acid C16:0 and C18:0 in stool and the decreased absorption coefficient of these two fatty acids.

Interactions of Divalent Cations

Complex interactions among cations are also apparent from the correlation analysis of calcium, phosphorus and magnesium excreted in stool, indicating calcium phosphate (fecal calcium and magnesium correlation: calcium with feedings: r=0.83) or magnesium phosphate (fecal phosphorus and magnesium correlation: calcium between feedings: r=0.78 and calcium with feedings: r=0.77) may be formed. Calcium-phosphorus-fat complex may also be formed as the correlations among fecal lipid, calcium, and phosphorus were quite significant.

Intraluminal calcium can modulate lipid dynamics of intestinal brush-border membrane in rat (22), and the change in membrane physiological processes may further complicate the mineral interactions. In vivo calcium gavage feeding results in a decrease in sphingomyelinase and an increase in sphingomyelin synthase in brush-border membrane, which enhance the synthesis of sphingomyelin in the membrane and, consequently,

decreases fluidity of brush-border membrane. The change in membrane lipid composition and fluidity may influence certain enzyme activities, receptor functions, permeability, and some ion channels within the membrane. It is possible that absorption of fatty acids, magnesium and phosphorus was affected in this manner.

4. Absorption Coefficients of Individual Fatty Acids

Fatty acids C12:0 and C14:0 were efficiently absorbed in the infants receiving calcium supplementation. However, fatty acids with sixteen and eighteen carbons seem to be affected by the calcium supplementation, among which C18:0 was most impaired (pooled absorption coefficient: 60.59%) because of the increased fecal loss of long-chain saturated fatty acids by forming insoluble calcium soaps in the intestinal lumen of the infants receiving calcium supplementation. This is in an agreement with other studies.

5. Fecal Lipid Fractions

The percentage distribution of fecal lipid in this study is contrary to that observed in other studies. It has been reported that the calcium salt fraction in stool of normal nonsupplemented newborn infants accounts for 25% of total fecal lipid; another 56% of fecal lipid is neutral lipid and the remainder is ionized fatty acids and bound lipid (8). Using the same methodology, the four fractions of fecal lipid in this study are distributed as follows: neutral lipid 54.4%, calcium salts 2.5%, free fatty acids 21.5% and compound lipid 21.7%. The difference in the calcium

salt fraction of this study cannot be readily explained. Perhaps, the influence of calcium supplementation on fat absorption is minimal and should not be done, or the mechanism of increased lipid excretion as suggested by the calcium-fat soap formation should be questioned.

Another study with either a lard-modified formula or an unmodified formula shows that greater than 90% of fecal lipid is free fatty acid (23) in newborn infants fed either formula. The high level of intestinal lipolysis was not seen in this study. Instead, free fatty acids in stool gradually decreased during the first four postnatal weeks while neutral lipid was increasing, indicating that the intestinal hydrolysis of milk lipid was progressively reduced at first weeks of life. A possible explanation is that the amount of lingual and gastric lipase is temporarily insufficient while its production and that of intestinal lipolytic enzymes are still quite limited at the early stage of life. The longitudinal change of these two fractions was more obvious in the infants receiving calcium between feedings, which might further indicate relatively less activity of lipase(s) at a low concentration of luminal calcium.

In conclusion, calcium and phosphorus content of preterm human milk decreased gradually in the first four weeks of life. Increased calcium intake resulted in the greater calcium loss from stool. Calcium supplementation between feedings further reduced the intestinal absorption of milk lipid, calcium, phosphorus, and magnesium compared to the supplementation with feedings. The absorption coefficients of C16:0 and C18:0 were most diminished in these infants, who also showed less weight

gain during the balance study. Therefore, it is not appropriate to feed calcium supplement between feedings to premature infants fed preterm human milk in the first four weeks of life. Yet intraluminal interactions of milk lipid, calcium, phosphorus, magnesium, and, possibly, other nutrients in human milk, as well as the role of calcium in the intestinal lipohydrolysis of milk, and fecal lipid fractionation need to be further elucidated.

E: REFERENCES:

1. Sauer P, Dane H, Visser H. New standards for neonatal thermal environment of health very low birth weight infants in week one of life. Arch Dis Child. 59: 18-22, 1984.

2. McIntyre. Flame photometry. Adv Clin Chem. 4: 1-28, 1961.

3. Fiske C, Subbarow Y. Calorimetric determination of phosphorus. J Bioch Chem. 66: 375-400, 1925.

4. Codex Alimentarius Commission: Methods of analysis for food for infants and children. In: Codex Alimentarius, Vol 9, Ed 1. Joint FAO/WHO Food Standards Program, Rome, 1982, Food and Agriculture Organization in the United Nations, pp.36.

5. Bitman J, Wood DL, Mehta NR, Hamosh P, Hamosh M. Lipolysis of triacylglycerols in human milk at low temperature: a note of caution. J Pediatr Gastroenterol Nutr. 2: 521-4, 1983.

6. Chappell J, Clandinin MT, Kearney-Volpe C, Reichman V, Swyer P. Fatty acid balance studies in premature infants fed human milk or formula: effect of calcium supplementation. J Pediatr. 108: 439-47, 1986.

7. Jeejeebhoy K, Ahmad S, Kozak S. Determination of fecal fats containing both medium and long chain triglycerides and fatty acids. Clin Biochem. 3: 157-63, 1970.

8. Watkins JB, Bliss CM, Donaldson RM, Lester R. Characterization of newborn fecal lipid. Pediatrics 53: 511-5, 1974.

9. Butte NF, Garza C, Johnson CA, Smith EO'B, Nichols BL. Longitudinal changes in milk composition of mothers delivering preterm and term infants. Early Hum Dev. 9: 153-62, 1984.

10. Atkinson S, Clandinin MT. Calcium supplementation of mother's milk for low birthweight infants: problems related to absorption and retention. Nutrition Res. 7: 813-23, 1987.

11. Chan GM. Human milk calcium and phosphate levels of mothers delivering term and preterm infants. J Pediatr Gastroenterol Nutr. 1 (2): 201-5, 1982.

12. Atkinson S, Clandinin MT. Calcium supplementation of mother's milk for low birthweight infants: problems related to absorption and retention. Nutrition Res. 7: 813-23, 1987.

13. Gross SJ, David RJ, Bauman L, Tomarelli RM. Nutritional composition of milk produced by mothers delivering preterm. J Pediatr. 96: 641-4, 1980.

14. Atkinson SA, Radde IC, Anderson GH. Macromineral balances in premature infants fed their own mothers' milk or formula. J Pediatr. 102 (1): 99-106, 1983.

15. Atkinson S, Bryan M, Anderson GH. Human milk feeding in premature infants: protein, fat and carbohydrate balances in the first two weeks of life. J Pediatr. 99: 617-24, 1981.

16. Chessex P, Reichman B, Verellen G, Putet G, Smith JM, Hein T, Swyer P. Quality of growth in premature infants fed their own mother's milk. J Pediatr. 102: 107-12, 1983.

17. Hamosh M. Lipid metabolism in premature infants. Biol Neonate 52 Suppl. 1: 50-64, 1987.

18. Borgstrom B. The importance of phospholipids, pancreatic phospholipase A2, fatty acid and other factors for the digestion of dietary fat: in vitro experiments with porcine enzymes. Gastroenterology 78: 954-62, 1980.

19. Borgstrom B. On the mechanism of pancreatic lipolysis of glycerides. Biochim Biophys Acta 13: 491-504, 1954.

20. Larsson A, Erlan-Albertsson C. The identity and properties of two forms of activated colipase from porcine pancrease. Biochim Biophys Acta 664: 538-48, 1981.

21. Alvarez FJ, Stella VJ. The role of calcium ions and bile salts on the pancreatic lipase-catalyzed hydrolysis of triglyceride emulsions stabilized with lecithin. Pharm Res. 6 (6): 449-57, 1989.

22. Dudeja PK, Brasitus TA, Dahiya R, Brown MD, Thomas D, Lau K. Intraluminal calcium modulates lipid dynamics of rat intestinal brush-border membranes. Am J Physiol. 252 (3 Pt 1): G398-403, 1987.

23. Verkade HJ, Hoving EB, Muskiet FAJ, Martini IA, Jansen G, Okken A, Vonk RJ, Bijleveld CMA. Fat absorption in neonates: comparisons of long-chain fatty acid and triglyceride compositions of formula, feces, and blood. Am J Clin Nutr. 53: 643-51, 1991.

Appendix I

EXTRACTION OF MILK LIPID

(Modified Folch Extraction)

1. Weigh approximately 0.2g freeze dried milk in a 50ml glass test tube with a screw cap and record the weight.
2. Add 1ml distilled deionized water and vortex.
3. Add 17.25ml Chloroform:Methanol solution (2:1 with Butylated Hydroxyl Toluene 0.005%) and 7.5mg internal standard Triglyceride C19:0 (0.75ml, 100mg/10ml in Chloroform:Methanol solution) in Fume Hood.
4. Polytron for 30 seconds.
5. Add 5ml 0.9% NaCl, vortex, and centrifuge at 2,000RPM for 10 minutes.
6. Extract 8ml chloroform carefully to a 15ml preweighed screw tube.
7. Evaporate to dryness with nitrogen protection in 50°C heating block.
8. Reweigh the tube, record the weight, and calculate the total weight of milk lipid: reweight of the tube (with dried lipid) - preweight of the blank tube.
9. Saponify the lipid by adding 2ml 0.5N methanolic KOH and heating at 100-105°C for 1 hour.
10. Methylate the sample by adding 2ml hexane and 1.5ml 14% Boron Trifluoride Methanol and heating at 100-105°C for 1 hour.
11. Add 2ml water, vortex, and centrifuge at 2,000RPM for 10 minutes.
12. Transfer the hexane to a 1.8ml vial, evaporate to dryness, and store in freezer (-20°C) under nitrogen protection for future GC analysis.

Appendix II

EXTRACTION OF FECAL LIPID

(Modified Method of Watkins and Bliss)

1. Weigh approximately 0.2mg freeze dried stool in a 50ml screw tube.
2. Add 5ml water and vortex for 2 minutes.
3. Add 10ml ether, vortex for 1 minute, and centrifuge at a maximum speed of 2,000RPM for 30 seconds.
4. Transfer the ether to a 15ml capped COREX tube and evaporate to 10ml at 40°C.
5. Repeat steps #3 an #4 another two times.

Ether Phase:

6. Centrifuge the COREX tube at 11,200RPM (i.e. 15,000G) at -5 to -10°C for 15 minutes to separate neutral lipid and calcium salts in ether.

Neutral lipid separation:

7. Transfer all neutral lipid-containing ether to a 25ml screw tube.
8. Add 10mg internal standard TG C19:0 (4ml, 2.5mg/1ml in hexane) and evaporate to dryness.
9. Saponify the lipid by adding 2ml 0.5N methanolic KOH and heating at 100-105°C for 1.5 hour.
10. Methylate the sample by adding 2ml hexane and 3ml 14% BF_3 and heating at 100-105°C for 1 hour.
11. Add 5ml water, vortex, and centrifuge at a maximum speed of 2,000RPM

for 30 seconds.

12. Transfer the hexane to a 1.8ml vial and evaporate to dryness.

Calcium salts separation:

13. Add 2ml water, 7-12 drops of 6N HCl and 2ml ether to the COREX tube and scrub the wall with Ca salt deposition repeatedly using a Pasteur pipet.
14. Transfer the ether-water to a 50ml screw tube.
15. Repeat steps #13 and #14 another two times.
16. Vortex and transfer ether to a 25ml screw tube and evaporate to dryness.
17. Repeat step #16 another two times.
18. Add 1mg internal standard FA C19:0 (2ml, 0.5mg/1ml in hexane).
19. Methylate the sample by adding 1ml 14% BF_3 and heating at 100-105°C for 1 hour.
20. Add 5ml water, vortex, and centrifuge at a maximum speed of 2,000RPM for 30 seconds.
21. Transfer the hexane to a 1.8ml vial and evaporate to dryness.

Water Phase:

22. Acidify the water phase to pH 1 by adding 6N HCl.
23. Add 20ml Chloroform:Methanol (2:1, 0.01% BHT) and 4ml saturated NaCl and vortex for 30 seconds.
24. Centrifuge at 2,000RPM for 5 minutes.

Free fatty acid separation:

25. Extract all chloroform to a 25ml screw tube and evaporate to dryness.
26. Add internal standard FA C19:0 4mg (2ml, 2mg/1ml in hexane) and 2ml 14% BF_3 and methylate at 100-105°C for 1 hour.
27. Add 5ml water, vortex, and centrifuge at a maximum speed of 2,000RPM for 30 seconds.
28. Transfer the hexane to a 1.8ml vial and evaporate to dryness.

Compound lipid separation:

29. Alkalize the remaining water phase to pH 10 by adding 10N NaOH.
30. Add internal standard TG C19:0 2mg (2ml, 1mg/1ml in hexane) and saponify it at 100-105°C overnight.
31. Reacidify to pH 1 by adding 6N HCl.
32. Add 20ml Chloroform:Methanol (2:1, 0.01% BHT) and 4ml saturated NaCl and vortex.
33. Centrifuge at 2,000RPM for 5 minutes.
34. Extract chloroform to a 25ml screw tube and evaporate to dryness.
35. Methylate the sample by adding 2ml hexane and 2ml 14% BF_3 and heating at 100-105°C for 1 hour.
36. Add 5ml water, vortex and centrifuge at a maximun speed of 2,000RPM for 30 seconds.
37. Transfer the hexane to a 1.8ml vial and evaporate to dryness.